

RISSO'S DOLPHIN (*Grampus griseus*): Western North Atlantic Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Risso's dolphins are distributed worldwide in tropical and temperate seas, and in the Northwest Atlantic occur from Florida to eastern Newfoundland (Leatherwood *et al.* 1976; Baird and Stacey 1990). Off the northeast U.S. coast, Risso's dolphins are distributed along the continental shelf edge from Cape Hatteras northward to Georges Bank during spring, summer, and autumn (CETAP 1982; Payne *et al.* 1984). In winter, the range is in the mid-Atlantic Bight and extends outward into oceanic waters (Payne *et al.* 1984). In general, the population occupies the mid-Atlantic continental shelf edge year round, and is rarely seen in the Gulf of Maine (Payne *et al.* 1984). During 1990, 1991 and 1993, spring/summer surveys conducted along the continental shelf edge and in deeper oceanic waters sighted Risso's dolphins associated with strong bathymetric features, Gulf Stream warm-core rings, and the Gulf Stream north wall (Waring *et al.* 1992; 1993; Hamazaki 2002). There is no information on stock structure of Risso's dolphin in the western North Atlantic, or to determine if separate stocks exist in the Gulf of Mexico and Atlantic. In 2006, a rehabilitated adult male Risso's dolphin stranded and released in the Gulf of Mexico off Florida was tracked via satellite to waters off Delaware (Wells *et al.* 2008). The Gulf of Mexico and Atlantic stocks are currently being treated as two separate stocks.

POPULATION SIZE

Total numbers of Risso's dolphins off the U.S. or Canadian Atlantic coast are unknown, although eight abundance estimates are available from selected regions for select time periods. Sightings were almost exclusively in the continental shelf edge and continental slope areas (Figure 1). The best abundance estimate for Risso's dolphins is the sum of the estimates from the two 2004 U.S. Atlantic surveys, 20,479 (CV=0.59), where the estimate from the northern U.S. Atlantic is 15,053 (CV=0.78), and from the southern U.S. Atlantic is 5,426 (CV=0.54). This joint estimate is considered best because these two surveys together have the most complete coverage of the population's habitat.

Earlier abundance estimates

Please see appendix IV for earlier abundance estimates. As recommended in the GAMMS Workshop Report (Wade and Angliss 1997), estimates older than eight years are deemed unreliable, therefore should not be used for PBR determinations. Further, due to changes in survey methodology these data should not be used to make comparisons to more current estimates.




Figure 1. Distribution of Risso's dolphin sightings from NEFSC and SEFSC shipboard and aerial surveys during the summers of 1998, 1999, 2002, 2004, 2006 and 2007. Isobaths are the 100-m, 1,000-m, and 4,000-m depth contours.

Recent surveys and abundance estimates

An abundance estimate of 9,311 (CV=0.76) Risso's dolphins was obtained from an aerial survey conducted in July and August 2002 which covered 7,465 km of trackline over waters from the 1,000 m depth contour on the southern edge of Georges Bank to Maine (Table 1; Palka 2006). The value of $g(0)$ used for this estimation was derived from the pooled data of 2002, 2004 and 2006 aerial survey data.

An abundance estimate of 15,054 (CV=0.78) Risso's dolphins was obtained from a line-transect sighting survey conducted during 12 June to 4 August 2004 by a ship and plane that surveyed 10,761 km of track line in waters north of Maryland (38°N) to the Bay of Fundy (45°N) (Table 1; Palka 2006). Shipboard data were collected using the two independent team line transect method and analyzed using the modified direct duplicate method (Palka 1995) accounting for biases due to school size and other potential covariates, reactive movements (Palka and Hammond 2001), and $g(0)$, the probability of detecting a group on the track line. Aerial data were collected using the Hiby circle-back line transect method (Hiby 1999) and analyzed accounting for $g(0)$ and biases due to school size and other potential covariates (Palka 2005).

A shipboard survey of the U.S. Atlantic outer continental shelf and continental slope (water depths >50 m) between Florida and Maryland (27.5 and 38°N latitude) was conducted during June-August 2004. The survey employed two independent visual teams searching with 25x bigeye binoculars. Survey effort was stratified to include increased effort along the continental shelf break and Gulf Stream front in the mid-Atlantic. The survey included 5,659 km of trackline, and recorded a total of 473 cetacean sightings. Sightings were most frequent in waters north of Cape Hatteras, North Carolina along the shelf break. Data were analyzed to correct for visibility bias ($g(0)$) and group-size bias employing line-transect distance analysis and the direct duplicate estimator (Palka 1995; Buckland *et al.* 2001). The resulting abundance estimate for Risso's dolphins between Florida and Maryland was 5,426 (CV =0.54).

An abundance estimate of 14,408 (CV=0.38) Risso's dolphins was obtained from an aerial survey conducted in August 2006 which covered 10,676 km of trackline in the region from the 2,000 m depth contour on the southern edge of Georges Bank to the upper Bay of Fundy and to the entrance of the Gulf of St. Lawrence (Table 1; Palka, pers. comm.). The value of $g(0)$ used for this estimation was derived from the pooled data of 2002, 2004 and 2006 aerial survey data.

Month/Year	Area	N_{best}	CV
Aug 2002	Georges Bank to Maine coast	9,311	0.76
Jun-Aug 2004	Maryland to Bay of Fundy	15,053	0.78
Jun-Aug 2004	Florida to Maryland	5,426	0.54
Jun-Aug 2004	Florida to Bay of Fundy (COMBINED)	20,479	0.59
Aug 2006	S. Gulf of Maine to upper Bay of Fundy to Gulf of St. Lawrence	14,408	0.38

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for Risso's dolphins is 20,479 (CV=0.59), obtained from the 2004 surveys. The minimum population estimate for the western North Atlantic Risso's dolphin is 12,920.

Current Population Trend

There are insufficient data to determine population trends for this species.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a “recovery” factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is 12,920. The maximum productivity rate is 0.04, the default value for cetaceans (Barlow *et al.* 1995). The “recovery” factor, which accounts for endangered, depleted, threatened stocks, or stocks of unknown status relative to optimum sustainable population (OSP) is assumed to be 0.48 because the CV of the average mortality estimate is between 0.3 and 0.6 (Wade and Angliss 1997). PBR for the western North Atlantic stock of Risso’s dolphin is 124.

ANNUAL HUMAN-CAUSED MORTALITY

Total annual estimated average fishery-related mortality or serious injury to this stock during 2003-2007 was 26 Risso’s dolphins (CV=0.32; Table 2).

Fishery Information

Detailed fishery information is reported in Appendix III.

Earlier Interactions

Prior to 1977, there was no documentation of marine mammal bycatch in distant-water fleet (DWF) activities off the northeast coast of the U.S. With implementation of the Fisheries Conservation and Management Act in that year, an observer program was established which recorded fishery data and information on incidental bycatch of marine mammals. NMFS foreign-fishery observers have reported four deaths of Risso’s dolphins incidental to squid and mackerel fishing activities in the continental shelf and continental slope waters between March 1977 and December 1991 (Waring *et al.* 1990; NMFS unpublished data).

In the pelagic drift gillnet fishery fifty-one Risso’s dolphin mortalities were observed between 1989 and 1998. One animal was entangled and released alive. Bycatch occurred during July, September and October along continental shelf edge canyons off the southern New England coast. Estimated annual mortality and serious injury (CV in parentheses) attributable to the drift gillnet fishery was 87 in 1989 (0.52), 144 in 1990 (0.46), 21 in 1991 (0.55), 31 in 1992 (0.27), 14 in 1993 (0.42), 1.5 in 1994 (0.16), 6 in 1995 (0), 0 in 1996, no fishery in 1997, 9 in 1998 (0).

In the pelagic pair trawl fishery, one mortality was observed in 1992. Estimated annual fishery-related mortality (CV in parentheses) attributable to the pelagic pair trawl fishery was 0.6 dolphins in 1991 (1.0), 4.3 in 1992 (0.76), 3.2 in 1993 (1.0), 0 in 1994 and 3.7 in 1995 (0.45).

Pelagic Longline

Pelagic longline bycatch estimates of Risso’s dolphins in 1998, 1999, and 2000 were obtained from Yeung (1999), Yeung *et al.* (2000), and Yeung (2001), respectively. Bycatch estimates for 2001 and 2002, 2003, 2004, 2005, 2006 and 2007 were obtained from Garrison (2003), Garrison and Richards (2004), Garrison (2005), Fairfield Walsh and Garrison (2006), Fairfield Walsh and Garrison (2007), and Fairfield and Garrison (2008). Most of the estimated marine mammal bycatch was from U.S. Atlantic EEZ waters between South Carolina and Cape Cod. Excluding the Gulf of Mexico, from 1992 to 2000 one mortality was observed in both 1994 and 2000, and 0 in other years. The observed numbers of seriously-injured but released alive individuals from 1992 to 2007 were, respectively, 2, 0, 6, 4, 1, 0, 1, 1, 1, 6, 4, 2, 2, 0, 0 and 1 (Cramer 1994; Scott and Brown 1997; Johnson *et al.* 1999; Yeung 1999; Yeung *et al.* 2000; Yeung 2001; Garrison 2003; Garrison and Richards 2004; Garrison 2005; Fairfield Walsh and Garrison 2006; Fairfield Walsh and Garrison 2007; Fairfield and Garrison 2008) (Table 2). Estimated annual fishery-related mortality (CV in parentheses) was 17 animals in 1994 (1.0), 41 in 2000 (1.0), 24 in 2001 (1.0), 20 in 2002 (0.86), and 0 in 2003 to 2007 (Table 2). Seriously injured and released alive animals were estimated to be 54 dolphins (0.7) in 1992, 0 in 1993, 120 (0.57) in 1994, 103 (0.68) in 1995, 99 (1.0) in 1996, 0 in 1997, 57 (1.0) in 1998, 22 (1.0) in 1999, 23 (1.0) in 2000, 45 (0.7) in 2001, 8 (1.0) in 2002, 40 (0.63) in 2003, 28 (0.72) in 2004, 3 (1.0), 0 in 2005, 0 in 2006, and 9 in 2007 (Table 2). The annual average combined mortality and serious injury for 2003-2007 is 16 Risso’s dolphins (CV =0.41; Table 2).

Northeast Sink Gillnet

Estimated annual mortalities (CV in parentheses) from this fishery are: 0 in 1999, 15 (1.06) in 2000, 0 in 2001-2004, 15 in 2005 (0.93), and 0 in 2006 and 2007 (Table 2). The 2003-2007 average mortality in this fishery is 3 Risso's dolphins (CV =0.93).

Mid-Atlantic Gillnet

A Risso's dolphin mortality was observed in this fishery for the first time in 2007. The resulting estimated annual mortality for 2007 was 33 (CV=0.73). The 2003-2007 average mortality in this fishery is 6 Risso's dolphins.

Table 2. Summary of the incidental mortality of Risso's dolphin (<i>Grampus griseus</i>) by commercial fishery including the years sampled (Years), the number of vessels active within the fishery (Vessels), the type of data used (Data Type), the annual observer coverage (Observer Coverage), the observed mortalities and serious injuries recorded by on-board observers, the estimated annual mortality and serious injury, the combined annual estimates of mortality and serious injury (Estimated Combined Mortality), the estimated CV of the combined estimates (Estimated CVs) and the mean of the combined estimates (CV in parentheses).											
Fishery	Years	Vessels ^b	Data Type ^a	Observer Coverage	Observed Serious Injury	Observed Mortality	Estimated Serious Injury	Estimated Mortality	Estimated Combined Mortality	Estimated CVs	Mean Annual Mortality
Pelagic Longline (excluding NED-E) ^c	03-07	63, 60, 60, 63, 62	Obs. Data Logbook	.09, .09, .06, .07, .08	2, 2, 0, 0, 1	0, 0, 0, 0, 0	40, 28, 3, 0, 9	0, 0, 0, 0, 0	40, 28, 3, 0, 9	.63, .72, 1, 0, .65	16(0.41)
Pelagic Longline - NED-E area only ^c	03	11	Obs. Data Logbook	1	0	1	0	1	1	1.0	1(1.0)
Northeast Sink Gillnet	03-07	1993=34 9 1998=30 1	Obs. Data Trip Logbook, Allocated Dealer Data	.03, .06, .07, .04, .07	0, 0, 0, 0, 0	0, 0, 1, 0, 0	0, 0, 0, 0, 0	0, 0, 15, 0, 0	0, 0, 15, 0, 0	0, 0, 0.93, 0, 0	3 (0.93)
Mid-Atlantic Gillnet	03-07	1993=34 9 1998=30 1	Obs. Data, Trip Logbook, Allocated Dealer Data	.01, .02, .03, .04, .04	0, 0, 0, 0, 0	0, 0, 0, 0, 1	0, 0, 0, 0, 0	0, 0, 0, 0, 33	0, 0, 0, 0, 33	0, 0, 0, 0, .73	6 (0.73)
TOTAL											26 (0.32)
^a	Observer data (Obs. Data) are used to measure bycatch rates and the data are collected within the Northeast Fisheries Observer Program. The Observer Program collects landings data (Weighout), and total landings are used as a measure of total effort for the coastal gillnet fishery.										
^b	Number of vessels in the fishery are based on vessels reporting effort to the pelagic longline logbook.										
^c	An experimental program to test effects of gear characteristics, environmental factors, and fishing practices on marine turtle bycatch rates in the Northeast Distant (NED-E) water component of the fishery was conducted from June 1, 2001-December 31, 2003. Observer coverage was 100% during this experimental fishery. Summaries are provided for the pelagic longline EXCLUDING the NED-E area in one row and for ONLY the NED in the second row (Garrison 2003; Garrison and Richards 2004) The NED area was reopened in June 2004, so 2004 - 2007 bycatch analysis includes this area.										

Other mortality

From 2003 to 2007, 75 Risso's dolphin strandings were recorded along the U.S. Atlantic coast (NMFS unpublished data). Six animals during this time period had indications of human interaction, three of which were fishery interactions. In eastern Canada, one Risso's dolphin stranding was reported on Sable Island, Nova Scotia from 1970-1998 (Lucas and Hooker 2000).

A Virginia Coastal Small Cetacean Unusual Mortality Event (UME) occurred along the coast of Virginia from 1 May to 31 July 2004, when 66 small cetaceans, including one Risso's dolphin, stranded mostly along the outer (eastern) coast of Virginia's barrier islands

A Mid-Atlantic Offshore Small Cetacean UME was declared when 33 small cetaceans stranded from Maryland to Georgia between July and September 2004. The species involved are generally found offshore and are not expected to strand along the coast. Three Risso's dolphins were involved in this UME.

STATE	2003	2004	2005	2006	2007	TOTALS
Maine	0	2	0	1	0	3
New Hampshire	0	0	0	0	0	0
Massachusetts ^{a,d}	10	4	8	1	3	26
Rhode Island	0	1	1	0	0	2
Connecticut	0	0	0	0	0	0
New York	0	3	4	1	0	8
New Jersey	0	0	5	0	2	7
Delaware	0	1	1	0	1	3
Maryland	2	1	2	1	0	6
Virginia ^b	1	1	4	1	1	8
North Carolina ^c	1	2	2	1	0	6
South Carolina	0	0	0	0	0	0
Georgia	0	0	0	0	0	0
Florida	1	3	0	0	1	5
EZ	0	1	0	0	0	1
TOTAL	15	19	27	6	8	75

a. One of the 2004 animals was mutilated, fluke cut off.
 b. One of the 2005 animals showed signs of fishery interaction.
 c. One of the 2006 animals showed signs of fishery interaction.
 d. 2003 includes 8 animals mass stranded in Massachusetts, 3 of which were released alive.

Stranding data probably underestimate the extent of fishery-related mortality and serious injury because all of the marine mammals that die or are seriously injured may not wash ashore, nor will all of those that do wash ashore necessarily show signs of entanglement or other fishery-interaction. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of fishery interaction.

STATUS OF STOCK

The status of Risso's dolphins relative to OSP in the U.S. Atlantic EEZ is unknown. The species is not listed as threatened or endangered under the Endangered Species Act. There are insufficient data to determine population trends for this species. The total U. S. fishery mortality and serious injury for this stock is not less than 10% of the calculated PBR and, therefore, can not be considered to be insignificant and approaching a zero mortality and serious injury rate. The 2003-2007 average annual human-related mortality does not exceed PBR; therefore, this is not a strategic stock.

REFERENCES CITED

- Baird, R.W. and P.J. Stacey 1990. Status of Risso's dolphin, *Grampus griseus*, in Canada. *Can. Field-Nat.* 105: 233-242.
- Barlow, J., S.L. Swartz, T.C. Eagle and P.R. Wade 1995. U.S. marine mammal stock assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6. 73 pp.
- Buckland, S.T., D.R. Anderson, K.P. Burnham, J.L. Laake, D.L. Borchers and L. Thomas 2001. Introduction to distance sampling: estimating abundance of biological populations. Oxford University Press. 432 pp.
- CETAP 1982. A characterization of marine mammals and turtles in the mid- and north Atlantic areas of the U.S. outer continental shelf, final report. Washington, DC, Bureau of Land Management. #AA551-CT8-48 538 pp.
- Cramer, J. 1994. Large pelagic logbook newsletter - 1993. NOAA Tech. Memo. NMFS-SEFSC-352. 19 pp.
- Fairfield, C.P. and L.P. Garrison 2008. Estimated bycatch of marine mammals and sea turtles in the US Atlantic pelagic longline fleet during 2007. NOAA Tech. Memo. NOAA NMFS-SEFSC-572. 62 pp.
- Fairfield Walsh, C. and L.P. Garrison 2006. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2005. NOAA Tech. Memo. NOAA NMFS-SEFSC-539. 52 pp.
- Fairfield Walsh, C. and L.P. Garrison 2007. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2006. NOAA Tech. Memo. NOAA NMFS-SEFSC-560. 54 pp.
- Garrison, L.P. 2003. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2001-2002. NOAA Tech. Memo. NMFS-SEFSC-515. 52 pp.
- Garrison, L.P. 2005. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2004. NOAA Tech. Memo. NMFS-SEFSC-531. 57 pp.
- Garrison, L.P. and P.M. Richards 2004. Estimated bycatch of marine mammals and turtles in the U.S. Atlantic pelagic longline fleet during 2003. NOAA Tech. Memo. NMFS-SEFSC-527. 57 pp.
- Hamazaki, T. 2002. Spatiotemporal prediction models of cetacean habitats in the mid-western North Atlantic Ocean (from Cape Hatteras, No. Carolina, USA to Nova Scotia, Canada). *Mar. Mamm. Sci.* 18(4): 920-939.
- Hiby, L. 1999. The objective identification of duplicate sightings in aerial survey for porpoise. Pages 179-189 *in*: G.W. Garner, S.C. Amstrup, J.L. Laake *et al.*, (eds.) *Marine Mammal Survey and Assessment Methods*. Balkema, Rotterdam.
- Johnson, D.R., C.A. Brown and C. Yeung 1999. Estimates of marine mammal and marine turtle catch by the U.S. Atlantic pelagic longline fleet in 1992-1997. NOAA Tech. Memo. NMFS- SEFSC-418. 70 pp.
- Leatherwood, S., D.K. Caldwell and H.E. Winn 1976. Whales, dolphins, and porpoises of the western North Atlantic. A guide to their identification. NOAA Tech. Rep. NMFS Circ. 396. 176 pp.
- Lucas, Z.N. and S.K. Hooker 2000. Cetacean strandings on Sable Island, Nova Scotia, 1970-1998. *Can. Field-Nat.* 114(1): 46-61.
- Palka, D. 1995. Abundance estimate of the Gulf of Maine harbor porpoise. *Rep. Int. Whal. Comm. (Special Issue)* 16: 27-50.
- Palka, D.L. 2005. Aerial surveys in the northwest Atlantic: estimation of $g(0)$, Proceedings of a Workshop on Estimation of $g(0)$ in Line-Transect Surveys of Cetaceans. European Cetacean Society's 18th Annual Conference; Kolmården, Sweden; Mar. 28, 2004.
- Palka, D.L. and P.S. Hammond 2001. Accounting for responsive movement in line transect estimates of abundance. *Can. J. Fish. Aquat. Sci* 58: 777-787.
- Payne, P.M., L.A. Selzer. and A.R. Knowlton. 1984. Distribution and density of cetaceans, marine turtles and seabirds in the shelf waters of the northeast U.S., June 1980 - Dec. 1983, based on shipboard observations. NMFS. NA81FAC00023 245 pp.
- Scott, G.P. and C.A. Brown 1997. Estimates of marine mammal and marine turtle catch by the U.S. Atlantic pelagic longline fleet in 1994-1995. M. L. Contribution. MIA-96/97-28.
- Wade, P.R. and R.P. Angliss 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Waring, G.T., C.P. Fairfield, C.M. Ruhsam and M. Sano 1992. Cetaceans associated with Gulf Stream Features off the Northeastern USA Shelf. *ICES [Int. Counc. Explor. Sea] C.M.* 1992/N:12.
- Waring, G.T., C.P. Fairfield, C.M. Ruhsam and M. Sano 1993. Sperm whales associated with Gulf Stream features off the northeastern USA shelf. *Fish. Oceanogr.* 2(2): 101-105.

- Waring, G.T., P. Gerrior, P.M. Payne, B.L. Parry and J.R. Nicolas 1990. Incidental take of marine mammals in foreign fishery activities off the northeast United States, 1977-1988. Fish. Bull. 88(2): 347-360.
- Wells, R.S., C.A. Manire, L. Byrd, D.R. Smith, J.G. Gannon, D. Fauquier and K.D. Mullin 2008. Movements and dive patterns of a rehabilitated Risso's dolphin, *Grampus griseus*, in the Gulf of Mexico and Atlantic Ocean. Mar. Mamm. Sci. 25(2): 420-429.
- Yeung, C. 1999. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1998. NOAA Tech. Memo. NMFS-SEFSC-430. 26 pp.
- Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1999-2000. NOAA Tech. Memo., NMFS, Southeast Fisheries Science Center, Miami, FL. NMFS-SEFSC-467. 43 pp.
- Yeung, C., S. Epperly and C.A. Brown 2000. Preliminary revised estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet, 1992-1999. NMFS, Miami Lab. PRD Contribution Number 99/00-13. 58 pp.