

STRIPED DOLPHIN (*Stenella coeruleoalba*): Hawaiian Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Striped dolphins are found in tropical to warm-temperate waters throughout the world (Perrin et al. 1994). There is an incongruity between the frequency of strandings and the infrequency of sightings of this species in Hawaii. Nitta (1991) found more stranding records of striped dolphins (13) than of any other species between 1936 and 1988, yet Shallenberger (1981) was aware of only two at-sea sightings, one near Niihau and one west of Oahu. A single sighting was made during recent systematic surveys within about 25 nmi of the main Hawaiian Islands (Figure 1). The Sea Life Park collecting crew never encountered striped dolphins from the early 1960s through the late 1970s, during their live-capture operations (Shallenberger 1981).

Striped dolphins have been intensively exploited in the western North Pacific, where three migratory stocks are provisionally recognized (Kishiro and Kasuya 1993). In the eastern Pacific all striped dolphins are provisionally considered to belong to a single stock (Dizon et al. 1994). For the Marine Mammal Protection Act (MMPA) stock assessment reports, striped dolphins within the Pacific U.S. Exclusive Economic Zone are divided into two discrete, non-contiguous areas: 1) waters off California, Oregon and Washington, and 2) waters around Hawaii (this report). Striped dolphins involved in eastern tropical Pacific tuna purse-seine fisheries are managed separately under the MMPA.




Figure 1. Location of the single sighting of striped dolphins during 1993-98 aerial surveys within about 25 nmi of the main Hawaiian Islands (see Appendix 2 for details on timing and location of survey effort). Outer line indicates approximate boundary of survey area.

POPULATION SIZE

Population estimates are available for Japanese waters (Miyashita 1993) and the eastern tropical Pacific (Wade and Gerrodette 1993), but it is not known whether any of these animals are part of the same population that occurs around the Hawaiian Islands. As part of the Marine Mammal Research Program of the Acoustic Thermometry of Ocean Climate (ATOC) study, a total of twelve aerial surveys were conducted within about 25 nmi of the main Hawaiian Islands in 1993, 1995 and 1998. An abundance estimate of 114 (CV=1.19) striped dolphins was recently calculated from the combined survey data (Mobley et al. 2000). This abundance underestimates the total number of striped dolphins within the U.S. EEZ off Hawaii, because areas around the Northwest Hawaiian Islands (NWHI) and beyond 25 nautical miles from the main islands were not surveyed.

Minimum Population Estimate

No data are available for a minimum population estimate. The log-normal 20th percentile of the combined 1993-98 abundance estimate is 52 striped dolphins. As with the best abundance estimate above, this includes only areas within about 25 nmi of the main Hawaiian Islands and is therefore an underestimate.

Current Population Trend

No data are available on current population trend.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

No data are available on current or maximum net productivity rate.

POTENTIAL BIOLOGICAL REMOVAL

The potential biological removal (PBR) level for this stock is calculated as the minimum population size (52)

times one half the default maximum net growth rate for cetaceans ($\frac{1}{2}$ of 4%) times a recovery factor of 0.50 (for a species of unknown status with no known fishery mortality; Wade and Angliss 1997), resulting in a PBR of 0.5 striped dolphins per year.

HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Fishery Information

No estimate of annual human-caused mortality and serious injury is available as there are no reports of direct or incidental takes of striped dolphins in Hawaiian waters (Nitta and Henderson 1993). However, mortality of other cetacean species has been observed in Hawaiian fisheries, and the gear types used in these fisheries are responsible for marine mammal mortality and serious injury in other fisheries throughout U.S. waters. Gillnets are used in Hawaiian waters and appear to capture marine mammals wherever they are used, and float lines from lobster traps and longlines can be expected to occasionally entangle whales (Perrin et al. 1994).

Interactions with cetaceans have been reported for all Hawaiian pelagic fisheries (Nitta and Henderson 1993), but no interactions with striped dolphins have been documented. None were observed hooked in the Hawaiian longline fishery between 1994 and 1998, with approximately 4.4% of all effort (measured as the number of hooks fished) observed (Kleiber 1999). Interaction rates between dolphins and the NWHI bottomfish fishery have been estimated based on studies conducted in 1990-1993, indicating that an average of 2.67 dolphin interactions, most likely involving bottlenose and rough-toothed dolphins, occurred for every 1000 fish brought on board (Kobayashi and Kawamoto 1995). Fishermen claim interactions with dolphins who steal bait and catch are increasing. It is not known whether these interactions result in serious injury or mortality of dolphins, nor whether striped dolphins are involved.

STATUS OF STOCK

The status of striped dolphins in Hawaiian waters relative to OSP is unknown, and there are insufficient data to evaluate trends in abundance. No habitat issues are known to be of concern for this species. They are not listed as “threatened” or “endangered” under the Endangered Species Act (1973), nor as “depleted” under the MMPA. Although information on striped dolphins in Hawaiian waters is limited, this stock would not be considered strategic under the 1994 amendments to the MMPA given the absence of reported fisheries related mortality. Insufficient information is available to determine whether the total fishery mortality and serious injury for striped dolphins is insignificant and approaching zero mortality and serious injury rate.

REFERENCES

- Dizon, A. E., W. F. Perrin, and P. A. Akin. 1994. Stocks of dolphins (*Stenella* spp. and *Delphinus delphis*) in the eastern tropical Pacific: a phylogeographic classification. NOAA Tech. Rep. NMFS 119, 20 pp.
- Kishiro, T. and T. Kasuya. 1993. Review of Japanese dolphin drive fisheries and their status. Rep. Int. Whal. Commn. 43:439-452.
- Kleiber, P. 1999. Estimates of marine mammal takes in the Hawaiian longline fishery. (Unpublished). Southwest Fisheries Science Center, NMFS, 2570 Dole St, Honolulu, HI, 96822-2396.
- Kobayashi, D. R. and K. E. Kawamoto. 1995. Evaluation of shark, dolphin, and monk seal interactions with Northwestern Hawaiian Island bottomfishing activity: a comparison of two time periods and an estimate of economic impacts. Fisheries Research 23: 11-22.
- Miyashita, T. 1993. Abundance of dolphin stocks in the western North Pacific taken by the Japanese drive fishery. Rep. Int. Whal. Commn. 43:417-437.
- Mobley, J. R., Jr, S. S. Spitz, K. A. Forney, R. A. Grotefendt, and P. H. Forestall. 2000. Distribution and abundance of odontocete species in Hawaiian waters: preliminary results of 1993-98 aerial surveys Admin. Rep. LJ-00-14C. Southwest Fisheries Science Center, National Marine Fisheries Service, P.O. Box 271, La Jolla, CA 92038. 26 pp.
- Nitta, E. 1991. The marine mammal stranding network for Hawaii: an overview. In: J.E. Reynolds III, D.K. Odell (eds.), Marine Mammal Strandings in the United States, pp.56-62. NOAA Tech. Rep. NMFS 98, 157 pp.
- Nitta, E. and J. R. Henderson. 1993. A review of interactions between Hawaii's fisheries and protected species. Mar. Fish. Rev. 55(2):83-92.
- Perrin, W. F., C. E. Wilson, and F. I. Archer, II. 1994. Striped dolphin *Stenella coeruleoalba* (Meyen, 1833). In: S. H. Ridgway and R. Harrison (eds.), Handbook of Marine Mammals, Vol.5: The First Book of Dolphins, pp 129-159. Academic Press, 416 pp.
- Perrin, W.F., G. P. Donovan and J. Barlow. 1994. Gillnets and Cetaceans. Rep. Int. Whal. Commn., Special Issue 15,

629 pp.

Shallenberger, E.W. 1981. The status of Hawaiian cetaceans. Final report to U.S. Marine Mammal Commission. MMC-77/23, 79pp.

Wade, P. R. and R. P. Angliss. 1997. Guidelines for Assessing Marine Mammal Stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U. S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12. 93 pp.

Wade, P. R. and T. Gerrodette. 1993. Estimates of cetacean abundance and distribution in the eastern tropical Pacific. Rep. Int. Whal. Commn. 43:477-493.