

HARBOR PORPOISE (*Phocoena phocoena*): Inland Washington Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

In the eastern North Pacific Ocean, the harbor porpoise ranges from Point Barrow, along the Alaskan coast, and down the west coast of North America to Point Conception, California (Gaskin 1984). Harbor porpoise primarily frequent coastal waters. Harbor porpoise are known to occur year-round in the inland trans-boundary area of Washington and British Columbia, Canada (Osborne et al. 1988), and along the Oregon/Washington coast (Barlow 1988, Barlow et al. 1988, Green et al. 1992). Aerial survey data from coastal Oregon and Washington, collected during all seasons, suggests that harbor porpoise distribution varies by depth (Green et al. 1992). Although distinct seasonal changes in abundance along the west coast have been noted, and attributed to possible shifts in distribution to deeper offshore waters during late winter (Dohl et al. 1983, Barlow 1988), harbor porpoise have also been conspicuously absent in offshore areas in late November (B. Taylor, pers. comm.) leaving a gap in the current understanding of their movements.

Stock discreteness in the eastern North Pacific was analyzed using mitochondrial DNA from samples collected along the west coast (Rosel 1992) and is summarized in Osmek et al. (1994). Two distinct mtDNA groupings or clades exist. One clade is present in California, Washington, British Columbia, and Alaska (no samples were available from Oregon), while the other is found only in California and Washington. Although these two clades are not geographically distinct by latitude, the results may indicate a low mixing rate for harbor porpoise along the west coast of North America. Investigation of pollutant loads in harbor porpoise ranging from California to the Canadian border also suggests restricted harbor porpoise movements (Calambokidis and Barlow 1991). Further genetic testing of the same data mentioned above, along with additional samples, found significant genetic differences for four of the six pair-wise comparisons between the four areas investigated: California, Washington, British Columbia, and Alaska (Rosel et al. 1995). These results demonstrate that harbor porpoise along the west coast of North America are not panmictic or migratory, and that movement is sufficiently restricted to evolve genetic differences. This is consistent with low movement suggested by genetic analysis of harbor porpoise specimens from the North Atlantic, where numerous stocks have been delineated with clinal differences over areas as small as the waters surrounding the British Isles.

Using the 1990-91 aerial survey data of Calambokidis et al. (1993) for water depths < 50 fathoms, Osmek et al. (1996) found significant differences in harbor porpoise mean densities ($z=5.9$, $p<0.01$) between the waters of coastal Oregon/Washington and inland Washington/southern British Columbia, Canada (i.e., Strait of Juan de Fuca/San Juan Islands). Although differences in density exist between coastal Oregon/Washington and inland Washington, a specific stock boundary line cannot be identified based upon biological or genetic differences. However, because harbor porpoise movements and rates of intermixing within the northeast Pacific are restricted, there has been a significant decline in harbor porpoise sightings within southern Puget Sound since the 1940s and, following a risk averse management strategy, two stocks are recognized to occur in Oregon and Washington waters (the Oregon/Washington Coast stock and the Inland Washington stock), with the boundary at Cape Flattery. Recent genetic evidence suggests that a population of animals at Spike Rock (on the northern coast of Washington, south of Cape Flattery) is more similar to the Inland Washington stock of harbor porpoise than to the Oregon/Washington Coast stock (S. Chivers, pers. comm.). All relevant data (e.g., additional genetic samples, contaminant studies, and satellite tagging) will be reviewed to determine whether to adjust the stock boundaries for harbor porpoise in Oregon and Washington waters.

In their assessment of California harbor porpoise, Barlow and Hanan (1995) recommended two stocks be recognized in California, with the stock boundary at the Russian River. Based on the above information four separate




Figure 1. Approximate distribution of harbor porpoise in the U.S. Pacific Northwest (shaded area). Stock boundaries separating the stocks are shown.

harbor porpoise stocks are recognized to occur along the west coast of the continental U.S. (see Fig. 1): 1) the Inland Washington stock, 2) the Oregon/Washington Coast stock, 3) the Northern California stock, and 4) the Central California stock. This report considers only the Inland Washington stock, with stock assessment reports for the Oregon/Washington Coast and both California stocks appearing in this volume. Three harbor porpoise stocks are also recognized in the inland and coastal waters of Alaska, including the Southeast Alaska, Gulf of Alaska, and Bering Sea stocks. The three Alaska harbor porpoise stocks are reported separately in the Stock Assessment Reports for the Alaska Region. The harbor porpoise occurring in British Columbia have not been included in any stock assessment report from either the Alaska Region or Pacific Northwest (Oregon/Washington).

POPULATION SIZE

Aerial surveys of the inside waters of Washington and southern British Columbia were conducted during August of 1996 (Calambokidis et al. 1997). These aerial surveys included the Strait of Juan de Fuca, San Juan Islands, Gulf Islands, and Strait of Georgia, which includes waters inhabited by harbor porpoise from British Columbia, as well as the Inland Washington stock. A total of 2,117 km of survey effort was completed within U.S. waters, resulting in an uncorrected abundance of 1,025 (CV=0.151) harbor porpoise in the inside waters of Washington (Calambokidis et al. 1997, Laake et al. 1997a). When corrected for availability and perception bias, using a correction factor of 3.42 ($1/g(0)$; $g(0)=0.292$, CV=0.366), the estimated abundance for the Inland Washington stock of harbor porpoise is 3,509 (CV=0.396) animals (Laake et al. 1997a, 1997b).

Minimum Population Estimate

The minimum population estimate (N_{MIN}) for this stock is calculated using Equation 1 from the PBR Guidelines (Wade and Angliss 1997): $N_{MIN} = N/\exp(0.842*\ln(1+[CV(N)]^2))^{1/2}$. Using the population estimate (N) of 3,509 and its associated CV(N) of 0.396, N_{MIN} for the Inland Washington stock of harbor porpoise is 2,545.

Current Population Trend

There are no reliable data on long-term population trends of harbor porpoise for most waters of Oregon, Washington, or British Columbia. For comparability to the 1996 survey, a re-analysis of the 1991 aerial survey data was conducted (Calambokidis et al. 1997). The abundance of harbor porpoise in the Inland Washington stock in 1996 was not significantly different than in 1991 (Laake et al. 1997a).

A different situation exists in southern Puget Sound where harbor porpoises are now rarely observed, a sharp contrast to 1942 when they were considered common in those waters (Scheffer and Slipp 1948). Although quantitative data for this area are lacking, marine mammal survey effort (Everitt et al. 1980), stranding records since the early 1970s (Osmek et al. 1995), and the results of harbor porpoise surveys of 1991 (Calambokidis et al. 1992) and 1994 (Osmek et al. 1995) indicate that harbor porpoise abundance has declined in southern Puget Sound. In 1994 a total of 769 km of vessel survey effort and 492 km of aerial survey effort conducted during favorable sighting conditions produced no sightings of harbor porpoise in southern Puget Sound. Reasons for the apparent decline are unknown, but it may be related to fishery interactions, pollutants, vessel traffic, or other activities that may affect harbor porpoise occurrence and distribution in this area (Osmek et al. 1995). Research to identify trends in harbor porpoise abundance is also needed for the other areas within inland Washington.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

A reliable estimate of the maximum net productivity rate is not currently available for harbor porpoise. Hence, until additional data become available, it is recommended that the cetacean maximum theoretical net productivity rate (R_{MAX}) of 4% (Wade and Angliss 1997) be employed for the Inland Washington harbor porpoise stock.

POTENTIAL BIOLOGICAL REMOVAL

Under the 1994 re-authorized Marine Mammal Protection Act (MMPA), the potential biological removal (PBR) is defined as the product of the minimum population estimate, one-half the maximum theoretical net productivity rate, and a recovery factor: $PBR = N_{MIN} \times 0.5R_{MAX} \times F_R$. The recovery factor (F_R) for this stock is 0.4, the value for a cetacean stock with an unknown population status and with a CV of mortality estimates greater than 0.8 (Wade and Angliss 1997). Thus, for the Inland Washington stock of harbor porpoise, $PBR = 20$ animals ($2,545 \times 0.02 \times 0.4$).

HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Fisheries Information

With the exception of 1994, NMFS observers monitored the northern Washington marine set gillnet fishery during 1992-97 (Gearin et al. 1994; P. Gearin, unpubl. data). For the entire area fished, observer coverage ranged from approximately 59 to 98% during those years. Fishing effort is conducted within the range of both harbor porpoise stocks (Oregon/Washington Coast and Inland Washington stocks) occurring in Washington State waters. Some of the animals taken in the inland waters portion of the fishery may have been animals from the coastal stock. Similarly, some of the animals taken in the coastal portion of the fishery (see stock assessment report for the Oregon/Washington Coast stock for details) may have been from the inland stock. For the purposes of this stock assessment report, the animals taken in the inland portion of the fishery are assumed to have belonged to the Inland Washington stock and the animals taken in the coastal portion of the fishery are assumed to have belonged to the Oregon/Washington Coast stock. Some movement of harbor porpoise between Washington's coastal and inland waters is likely, but it is currently not possible to quantify the extent of such movements. Accordingly, Table 1 includes data only from that portion of the northern Washington marine set gillnet fishery occurring within the range of the Inland Washington stock (those waters east of Cape Flattery), where observer coverage ranged from 6-80% between 1992 and 1997. Data from 1992-97 are included in Table 1, although the mean estimated annual mortality is calculated using the most recent 5 years of available data. No mortalities were observed in the inland portion of the fishery between 1992 and 1997. As noted above, there was no observer program in 1994. Little effort occurred in 1995 and observer coverage was lower than usual (24%). Effort increased in 1996, however, the observer coverage decreased to a low of 6%. In 1997, observer coverage increased to 80%, although little effort occurred. The mean estimated mortality for this fishery is zero harbor porpoise per year from this stock.

In 1993, as a pilot for future observer programs, NMFS in conjunction with the Washington Department of Fish and Wildlife (WDF&W) monitored all non-treaty components of the Washington Puget Sound Region salmon gillnet fishery (Pierce et al. 1994). Observer coverage was 1.3% overall, ranging from 0.9% to 7.3% for the various components of the fishery. No harbor porpoise mortalities were reported (Table 1). Pierce et al. (1994) cautioned against extrapolating these mortalities to the entire Puget Sound fishery due to the low observer coverage and potential biases inherent in the data. The area 7/7A sockeye landings represented the majority of the non-treaty salmon landings in 1993, approximately 67%. Results of this pilot study were used to design the 1994 observer programs discussed below.

In 1994, NMFS in conjunction with WDF&W conducted an observer program during the Puget Sound non-treaty chum salmon gillnet fishery (areas 10/11 and 12/12B). A total of 230 sets were observed during 54 boat trips, representing approximately 11% observer coverage of the 500 fishing boat trips comprising the total effort in this fishery as estimated from fish ticket landings (Erstad et al. 1996). No harbor porpoise were reported within 100 meters of observed gillnets. The Puget Sound treaty chum salmon gillnet fishery in Hood Canal (areas 12, 12B, and 12C) and Puget Sound treaty sockeye/chum gillnet fishery in the Strait of Juan de Fuca (areas 4B, 5, and 6C) were also monitored in 1994 (NWIFC 1995). No harbor porpoise mortalities were reported in the observer programs covering these treaty salmon gillnet fisheries, where observer coverage was estimated at 2.2% (based on % of total catch observed) and approximately 7.5% (based on % of observed trips to total landings), respectively.

Also in 1994, NMFS in conjunction with the WDF&W and the Tribes conducted an observer program to examine seabird and marine mammal interactions with the Puget Sound treaty and non-treaty sockeye salmon gillnet fishery (areas 7 and 7A). During this fishery, observers monitored 2,205 sets, representing approximately 7% of the estimated 33,086 sets occurring in the fishery (Pierce et al. 1996). There was one observed harbor porpoise mortality (one other was entangled and released alive with no indication the animal was injured), resulting in a mortality rate of 0.00045 harbor porpoise per set, which extrapolates to 15 mortalities (CV=1.0) for the entire fishery. In 1996, Washington Sea Grant Program conducted a test fishery in the non-treaty sockeye salmon gillnet fishery (area 7) to compare entanglement rates of seabirds and marine mammals and catch rates of salmon using three experimental gears and a control (monofilament mesh net). The experimental nets incorporated highly visible mesh in the upper quarter (50 mesh gear) or upper eighth (20 mesh gear) of the net or had low-frequency sound emitters attached to the corkline (Melvin et al. 1997). In 642 sets during 17 vessel trips, 2 harbor porpoise were killed in the 50 mesh gear.

Combining the estimates from the 1994 observer programs (15) with the northern Washington marine set gillnet fishery (0) results in an estimated mean mortality rate in observed fisheries of 15 harbor porpoise per year from this stock. It should be noted that the 1994 observer programs did not sample all segments of the entire Washington Puget Sound Region salmon set/drift gillnet fishery, and further, the extrapolation of total kill did not include effort for the unobserved segments of this fishery. Therefore, 15 is an underestimate of the harbor porpoise mortality due to the entire fishery. Though it is not possible to quantify what percentage of the Washington Puget Sound Region salmon set/drift gillnet

fishery was actually observed in 1994, the observer programs covered those segments of the fishery which had the highest salmon catches, the majority of vessel participation, and the highest likelihood of interaction with harbor porpoise (J. Scordino, pers. comm.). Accordingly, the estimated harbor porpoise mortality (15) appears to be only a slight underestimate for the fishery. See Appendix 1 of Barlow et al. (1997) for additional information, including a map depicting fishing areas, regarding the Washington Puget Sound Region salmon set/drift gillnet fishery.

Table 1. Summary of incidental mortality of harbor porpoise (Inland Washington stock) due to commercial fisheries from 1992 through 1997 and calculation of the mean annual mortality rate. Only data from 1993 to 1997 (or the most recent 5 years of available data) are used to calculate mean annual mortality (n/a indicates that data are not available).

Fishery name	Years	Data type	Range of observer coverage	Observed mortality (in given yrs.)	Estimated mortality (in given yrs.)	1993-97 Mean annual mortality
Northern WA marine set gillnet (inland waters)	92-97	obs data	6-80%	0, 0, n/a, 0, 0, 0	0, 0, n/a, 0, 0, 0	0
WA Puget Sound Region salmon set/drift gillnet (observer programs listed below covered segments of this fishery):	-	-	-	-	-	-
Puget Sound non-treaty salmon gillnet (all areas and species)	93	obs data	1.3%	0	0	see text
Puget Sound non-treaty chum salmon gillnet (areas 10/11 and 12/12B)	94	obs data	11%	0	0	0
Puget Sound treaty chum salmon gillnet (areas 12, 12B, and 12C)	94	obs data	2.2%	0	0	0
Puget Sound treaty chum and sockeye salmon gillnet (areas 4B, 5, and 6C)	94	obs data	7.5%	0	0	0
Puget Sound treaty and non-treaty sockeye salmon gillnet (areas 7 and 7A)	94	obs data	7%	1	15	15 (CV=1.0)
Observer program total						15 (CV=1.0)
				Reported mortalities		
WA Puget Sound Region salmon set/drift gillnet	90-97	self reports	n/a	6, 4, 6, 2, n/a, n/a, n/a, n/a	n/a	see text
Minimum total annual mortality						\$15 (CV=1.0)

An additional source of information on the number of harbor porpoises killed or injured incidental to commercial fishery operations is the self-reported fisheries information required of vessel operators by the MMPA. Self-reported fishery data from 1990 to 1997 for the Washington Puget Sound Region salmon set and drift gillnet fishery are shown in Table 1. Unlike the 1994 observer program data, the self-reported fisheries data cover the entire fishery. However, because logbook records (fisher self-reports required during 1990-94) are most likely negatively biased (Credle et al. 1994), these are considered to be minimum estimates of harbor porpoise mortality. Self-reported fisheries data are incomplete for 1994, not available for 1995, and considered unreliable after 1995 (see Appendix 4 of Hill and DeMaster 1998). Though the 1994 observer program data may underestimate the total fishery mortality for this stock, it is considered more reliable than the self-reported data. Thus, the self-reported fisheries data were not used in the mortality rate calculation.

Strandings of harbor porpoise wrapped in fishing gear or with injuries caused by interactions with gear are a final source of fishery-related mortality information. During the period from 1992 to 1997 the only reported fishery-related strandings of harbor porpoise occurred in 1992 (1 animal) and 1993 (1 animal). The mortalities likely occurred in the Washington Puget Sound Region salmon set and drift gillnet fishery. As the 1994 observer program already accounts for 15 harbor porpoise mortalities per year from this fishery, these strandings have not been included in Table 1.

There are few data concerning the mortality of marine mammals incidental to commercial gillnet fisheries in Canadian waters, which have not been monitored but are known to have taken harbor porpoise in the past (Barlow et al. 1994, Stacey et al. 1997). As a result, the number of harbor porpoise from this stock currently taken in the waters of southern British Columbia is not known.

STATUS OF STOCK

Harbor porpoise are not listed as “depleted” under the MMPA or listed as “threatened “ or “endangered” under the Endangered Species Act. Based on currently available data, the level of human-caused mortality and serious injury (15) is not known to exceed the PBR (20). Therefore, the Inland Washington harbor porpoise stock is not classified as strategic. The minimum total fishery mortality and serious injury for this stock (15) exceeds 10% of the calculated PBR (2.0) and, therefore, cannot be considered to be insignificant and approaching zero mortality and serious injury rate. The status of this stock relative to its Optimum Sustainable Population and population trends is unknown, although harbor porpoise sightings in the southern Puget Sound have declined since the 1940s.

Although this stock is not recognized as strategic at this time, there is cause for concern due to the following issues: 1) the estimated take level is close to exceeding the PBR (i.e., one additional observed mortality or serious injury in the area 7/7A sockeye drift gillnet fishery would increase the estimated annual take level above the PBR), 2) the extent to which harbor porpoise from U. S. waters frequent the waters of British Columbia, and are therefore subject to fishery-related mortality, is unknown, and 3) the mortality rate is based on observer data from a subset of the Washington Puget Sound Region salmon set and gillnet fishery.

REFERENCES

- Barlow, J. 1988. Harbor porpoise, *Phocoena phocoena*, abundance estimation for California, Oregon and Washington: I. Ship surveys. Fish. Bull. 86:417-432.
- Barlow, J., and D. Hanan. 1995. An assessment of the status of harbor porpoise in central California. Rep. Int. Whal. Commn. Special Issue 16:123-140.
- Barlow, J., C. W. Oliver, T. D. Jackson, and B. L. Taylor. 1988. Harbor porpoise, *Phocoena phocoena*, abundance estimation for California, Oregon and Washington: II. Aerial surveys. Fish. Bull. 86:433-444.
- Barlow, J., R. W. Baird, J. E. Heyning, K. Wynne, A. M. Manville, II, L. F. Lowry, D. Hanan, J. Sease, and V. N. Burkanov. 1994. A review of cetacean and pinniped mortality in coastal fisheries along the west coast of the USA and Canada and the east coast of the Russian Federation. Rep. Int. Whal. Commn. Special Issue 15:405-425.
- Barlow, J., K. A. Forney, P. S. Hill, R. L. Brownell, Jr., J. V. Carretta, D. P. DeMaster, F. Julian, M. S. Lowry, T. Ragen, and R. R. Reeves. 1997. U.S. Pacific marine mammal stock assessments: 1996. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SWFSC-248. 223 pp.
- Calambokidis, J., and J. Barlow. 1991. Chlorinated hydrocarbon concentrations and their use for describing population discreteness in harbor porpoises from Washington, Oregon, and California. Reynolds, J. E., III, and D. K. Odell (eds.), Proceedings of the Second Marine Mammal Stranding Workshop: 3-5 December 1987, Miami, Florida. NMFS, NOAA Technical Rep. NMFS 98:101-110.
- Calambokidis, J., J. R. Evenson, J. C. Cabbage, P. J. Gearin, and S. D. Osmek. 1992. Harbor porpoise distribution and

- abundance off Oregon and Washington from aerial surveys in 1991. Final Rep. by Cascadia Research, Olympia, WA, to National Marine Mammal Laboratory, AFSC, NMFS, Seattle, WA. 44 pp.
- Calambokidis, J., J. C. Cubbage, J. R. Evenson, S. D. Osmeck, J. L. Laake, P. J. Gearin, B. J. Turnock, S. J. Jeffries, and R. F. Brown. 1993. Abundance estimates of harbor porpoise in Washington and Oregon waters. Final Rep. by Cascadia Research, Olympia, WA, to National Marine Mammal Laboratory, AFSC, NMFS, Seattle, WA. 55 pp.
- Calambokidis, J., S. D. Osmeck, and J. L. Laake. 1997. Aerial surveys for marine mammals in Washington and British Columbia inside waters. Final Rep. by Cascadia Research, Olympia, WA, to National Marine Mammal Laboratory, AFSC, NMFS, Seattle, WA. 96 pp.
- Chivers, S. Southwest Fisheries Science Center, NMFS, P.O. Box 271, La Jolla, CA 92038.
- Credle, V. R., D. P. DeMaster, M. M. Merklein, M. B. Hanson, W. A. Karp, and S. M. Fitzgerald (eds.). 1994. NMFS observer programs: minutes and recommendations from a workshop held in Galveston, Texas, November 10-11, 1993. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-94-1. 96 pp.
- Dohl, T. P., R. C. Guess, M. L. Duman, and R. C. Helm. 1983. Cetaceans of central and northern California, 1980-1983: Status, abundance, and distribution. OCS study MMS 84-0045. Pacific OCS Region Minerals Management Service, 1340 Sixth Street, Los Angeles, CA 90014. 284 pp.
- Erstad, P., S. L. Jeffries, and D. J. Pierce. 1996. 1994 Report for the Puget Sound fishery observer program in management areas 10/11 & 12/12B: nontreaty chum gill net fishery. Wash. Dept. Fish and Wildlife., Olympia, WA. 14 pp.
- Everitt, R. D., C. H. Fiscus, R. L. DeLong. 1980. Northern Puget Sound marine mammals. U.S. Department of Commerce EPA-6000/7-80-139. 134 pp.
- Gaskin, D. E. 1984. The harbour porpoise *Phocoena phocoena* (L.): Regional populations, status, and information on direct and indirect catches. Rep. Int. Whal. Commn. 34:569-586.
- Gearin, P. J. National Marine Mammal Laboratory, AFSC, NMFS, 7600 Sand Point Way NE, Seattle, WA 98115.
- Gearin, P. J., S. R. Melin, R. L. DeLong, H. Kajimura, and M. A. Johnson. 1994. Harbor porpoise interactions with a chinook salmon set-net fishery in Washington state. Rep. Int. Whal. Commn. Special Issue 15:427-438.
- Green, G. A., J. J. Brueggeman, R. A. Grotefendt, C. E. Bowlby, M. L. Bonnel, and K. C. Balcomb. 1992. Cetacean distribution and abundance off Oregon and Washington, 1989-1990. Ch. 1, *In*: Brueggeman, J. J. (ed.), Oregon and Washington Marine Mammal and Seabird Surveys. Final Rep. OCS Study MMS 91-0093.
- Hill, P. S., and D. P. DeMaster. 1998. Alaska marine mammal stock assessments, 1998. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-97. 166 pp.
- Laake, J. L., R. L. DeLong, J. Calambokidis, and S. D. Osmeck. 1997a. Abundance and distribution of marine mammals in Washington and British Columbia inside waters, 1996. Annual rep. to the MMPA Assessment Program, Office of Protected Resources, NMFS, NOAA, 1335 East-West Highway, Silver Spring, MD 20910. 7 pp. Available at National Marine Mammal Laboratory, 7600 Sand Point Way NE, Seattle, WA 98115.
- Laake, J. L., J. Calambokidis, S. D. Osmeck, and D. J. Rugh. 1997b. Probability of detecting harbor porpoise from aerial surveys: Estimating $g(0)$. J. Wildl. Manage. 61(1):63-75.
- Melvin, E. F., L. L. Conquest, and J. K. Parrish. 1997. Seabird bycatch reduction: New tools for Puget Sound drift gillnet salmon fisheries. 1996 Sockeye and 1995 Chum Salmon Test Fisheries Final Report. Washington Sea Grant Program, Seattle, WA. 48pp.
- Northwest Indian Fisheries Commission. 1995. Monitoring of marbled murrelet and marine mammal interactions with 1994 tribal gillnet fisheries in northern Puget Sound, Hood Canal, and the Strait of Juan de Fuca. Final Report to NMFS Contract No. 52ABNF400087. Unpubl. report. Northwest Indian Fisheries Commission. 43 pp. Available at NWIFC, 6730 Martin Way E, Olympia, WA 98516.
- Osborne, R., J. Calambokidis, and E. M. Dorsey. 1988. A Guide to Marine Mammals of Greater Puget Sound. Island Publishers, Anacortes, WA. 191 pp.
- Osmeck, S. D., P. E. Rosel, A. E. Dizon, and R. L. DeLong. 1994. Harbor porpoise *Phocoena phocoena* population assessment studies for Oregon and Washington in 1993. Annual Rep. to the MMPA Assessment Program, Office of Protected Resources, NMFS, NOAA, 1335 East-West Highway, Silver Spring, MD 20910. 14 pp. Available at National Marine Mammal Laboratory, 7600 Sand Point Way NE, Seattle, WA 98115.
- Osmeck, S. D., M. B. Hanson, J. L. Laake, S. J. Jeffries, and R. L. DeLong. 1995. Harbor porpoise *Phocoena phocoena* population assessment studies for Oregon and Washington in 1994. Annual Rep. to the MMPA Assessment Program, Office of Protected Resources, NMFS, NOAA, 1335 East-West Highway, Silver Spring, MD 20910. 32 pp. Available at National Marine Mammal Laboratory, 7600 Sand Point Way NE, Seattle, WA 98115.

- Osmek, S., J. Calambokidis, J. Laake, P. Gearin, R. DeLong, J. Scordino, S. Jeffries, and R. Brown. 1996. Assessment of the status of harbor porpoises, *Phocoena phocoena*, in Oregon and Washington waters. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-76. 46 pp.
- Pierce, D. J., W. P. Ritchie, and R. Kreuziger. 1994. Preliminary findings of seabird interactions with the non-treaty salmon gill net fishery: Puget Sound and Hood Canal Washington. Unpubl. report. Washington Department of Fish and Wildlife, Olympia, WA. 39 pp. Available at WDFW, 600 Capitol Way N, Olympia, WA, 98501.
- Pierce, D. J., M. Alexandersdottir, S. J. Jeffries, P. Erstad, W. Beattie, and A. Chapman. 1996. Interactions of marbled murrelets and marine mammals with the 1994 Puget Sound sockeye gill net fishery. Final Report, Wash. Dept. Fish and Wildlife, Olympia, WA. 21 pp.
- Rosel, P. E. 1992. Genetic population structure and systematic relationships of some small cetaceans inferred from mitochondrial DNA sequence variation. Ph.D. Dissertation, Univ. Calif. San Diego, La Jolla, CA. 191 pp.
- Rosel, P. E., A. E. Dizon, and M. G. Haygood. 1995. Variability of the mitochondrial control region in populations of the harbour porpoise, *Phocoena phocoena*, on inter-oceanic and regional scales. Can. J. Fish. and Aquat. Sci. 52:1210-1219.
- Scheffer, V. B., and J. W. Slipp. 1948. The whales and dolphins of Washington state with a key to the cetaceans of the west coast of North America. Am. Midl. Nat. 39(2):257-337.
- Scordino, J. National Marine Fisheries Service, Northwest Region, 7600 Sand Point Way NE, Seattle, WA 98115.
- Stacey, P. J., R. W. Baird, and D. A. Duffus. 1997. A preliminary evaluation of incidental mortality of small cetaceans, in coastal fisheries in British Columbia, Canada. Mar. Mamm. Sci. 13:321-326.
- Taylor, B. Southwest Fisheries Science Center, NMFS, P. O. Box 271, La Jolla, CA 92038.
- Wade, P. R., and R. Angliss. 1997. Guidelines for assessing marine mammal stocks: report of the GAMMS workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12. 93 pp.