

RINGED SEAL (*Phoca hispida hispida*): Alaska Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Ringed seals have a circumpolar distribution and are found in all seasonally ice-covered seas of the Northern Hemisphere as well as in certain freshwater lakes (King 1983). Most taxonomists currently recognize five subspecies of ringed seals: *Phoca hispida hispida* in the Arctic Ocean and Bering Sea; *Phoca hispida ochotensis* in the Sea of Okhotsk and northern Sea of Japan; *Phoca hispida botnica* in the northern Baltic Sea; *Phoca hispida lagodensis* in Lake Ladoga, Russia; and *Phoca hispida saimensis* in Lake Saimaa, Finland. The lake-inhabiting subspecies are genetically isolated and those in the Baltic Sea and the Sea of Okhotsk apparently exchange genes with the Arctic subspecies minimally or not at all (Palo et al. 2001, Palo 2003, Kelly et al. 2009). The genetic structuring of *Phoca hispida hispida*, however, remains unresolved, and it may prove to be composed of multiple distinct populations (Kelly et al. 2010a). For the purposes of this stock assessment, the Alaska stock of ringed seals is considered the portion of *Phoca hispida hispida* that occurs within the U.S. Exclusive Economic Zone of the Beaufort, Chukchi, and Bering Seas (Fig. 12).



Figure 12. Approximate distribution of ringed seals (shaded area). The combined summer and winter distribution are depicted.

Throughout their range, ringed seals have an affinity for ice-covered waters and are well adapted to occupying both shorefast and pack ice (Kelly 1988a). They remain in contact with ice most of the year and use it as a platform for pupping and nursing in late winter to early spring, for molting in late spring to early summer, and for resting at other times of the year, although land haulouts may be increasingly used because of increases in summer sea ice retreat. In Alaskan waters, during winter and early spring when sea ice is at its maximal extent, ringed seals are abundant in the northern Bering Sea, Norton and Kotzebue Sounds, and throughout the Chukchi and Beaufort Seas. They occur as far south as Bristol Bay in years of extensive ice coverage but generally are not abundant south of Norton Sound except in nearshore areas (Frost 1985). Although details of their seasonal movements have not been adequately documented, it is thought that most ringed seals that winter in the Bering and Chukchi Seas migrate north in spring as the seasonal ice melts and retreats (Burns 1970) and have historically spent summer in the pack ice of the northern Chukchi and Beaufort seas, as well as in nearshore ice remnants in the Beaufort Sea (Frost 1985); however, very little pack ice exists in the summer months in the northern Chukchi and Beaufort seas. During summer, ringed seals range hundreds to thousands of kilometers to forage along ice edges or in highly productive open-water areas (Freitas et al. 2008, Kelly et al. 2010b). With the onset of freeze-up in the fall, ringed seal movements become increasingly restricted and seals that have summered in the Beaufort Sea are thought to move west and south with the advancing ice pack, with many seals dispersing throughout the Chukchi and Bering Seas while some remain in the Beaufort Sea (Frost and Lowry 1984, Crawford et al. 2012). Many adult ringed seals return to the same small home ranges they occupied during the previous winter (Kelly et al. 2010b).

POPULATION SIZE

Ringed seal population surveys in Alaska have used various methods and assumptions, had incomplete coverage of their habitats and range, and were conducted more than a decade ago; therefore, current, comprehensive, and reliable abundance estimates or trends for the Alaska stock are not available. Burns and Harbo (1972) conducted aerial surveys along the North Slope of Alaska (between Point Lay and Kaktovik) during June 1970, and reported a minimal estimate of 11,612 ringed seals in areas of shorefast ice. Frost and Lowry (1984) produced a rough estimate

of 40,000 ringed seals in the Alaskan Beaufort Sea during winter and spring by applying an assumed correction factor for availability bias (i.e., for seals not hauled out at the time of the surveys) to the average density observed from 7 years of aerial surveys in the Alaskan and Yukon Beaufort Sea and extrapolating over the entire area of the continental shelf. Their estimate during summer of 80,000 ringed seals was based on the assumption that this population doubles as seals from the Bering and Chukchi Seas move in with the receding ice edge. Based on an analysis of surveys conducted during the 1970s, Frost (1985) estimated 1 to 1.5 million ringed seals in Alaskan waters, of which 250,000 were estimated in shorefast ice. These estimates were considered conservative when compared with polar bear predation rates (Frost 1985); however, details of the analysis were not published. Frost et al. (1988) reported detailed methods and results of surveys conducted in the Alaskan Chukchi and Beaufort Seas during May-June 1985-1987. Survey effort was directed towards shorefast ice within 20 nmi of shore, though some areas of adjacent pack ice were also surveyed, and estimates were based on observed densities extrapolated over estimates of available habitat without correcting for availability bias. In the Chukchi Sea, total numbers of hauled out ringed seals in shorefast ice ranged from $18,400 \pm 1,700$ in 1985 to $35,000 \pm 3,000$ in 1986. The 1987 estimate of $20,200 \pm 2,300$ was similar to 1985. In the Beaufort Sea, the estimated number of ringed seals hauled out within the 20-m depth contour ranged from $9,800 \pm 1,800$ in 1985 to $13,000 \pm 1,600$ in 1986. The 1987 estimate ($19,400 \pm 3,700$) was considerably higher but may have included seals that had moved in from other areas as the ice began to break up (Frost et al. 1988). Frost et al. (2004) conducted surveys within 40 km of shore in the Alaskan Beaufort Sea during May-June 1996-1999, and observed ringed seal densities ranging from 0.81 seals/km² in 1996 to 1.17 seals/km² in 1999. Moulton et al. (2002) conducted similar, concurrent surveys in the Alaskan Beaufort Sea during 1997-1999 but reported substantially lower ringed seal densities than Frost et al. (2004). The reason for this disparity was unclear (Frost et al. 2004). Bengtson et al. (2005) conducted surveys in the Alaskan Chukchi Sea during May-June 1999 and 2000. While the surveys were focused on the coastal zone within 37 km of shore, additional survey lines were flown up to 185 km offshore. Population estimates were derived from observed densities corrected for availability bias using a haul-out model from 6 tagged seals. Ringed seal abundance estimates for the entire survey area were 252,488 (SE = 47,204) in 1999 and 208,857 (SE = 25,502) in 2000. The estimates from 1999 and 2000 in the Chukchi Sea only covered a portion of this stock's range and were conducted over a decade ago. Using the more recent estimates from surveys by Bengtson et al. (2005) and Frost et al. (2004) in the late 1990s, the total population in the Alaskan Chukchi and Beaufort Sea regions is at least 300,000 seals, which is likely an underestimate (Kelly et al. 2010a). Using the most recent estimates from surveys by Bengtson et al. (2005) and Frost et al. (2004) in the late 1990s and 2000, for purposes of the ESA status review of the species, Kelly et al. (2010a) estimated the total population in the Alaskan Chukchi and Beaufort Seas as at least 300,000 ringed seals, which Kelly et al. (2010a) state is likely an underestimate since the Beaufort surveys were limited to within 40 km of shore.

Minimum Population Estimate

The estimate of 300,000 seals presented in Kelly et al. (2010a) is based on estimates from surveys by Bengtson et al. (2005) and Frost et al. (2004) in the late 1990s and 2000. This is not considered a reliable abundance estimate, is likely an underestimate, and is based on surveys of a portion of the range and are greater than 8 years old. A reliable estimate of N_{MIN} for the total population in the Alaskan Chukchi and Beaufort Sea regions is not available.

Current Population Trend

Frost et al. (2002) reported that trend analysis based on an ANOVA comparison of observed seal densities in the central Beaufort Sea suggested marginally significant but substantial declines of 50% on shorefast ice and 31% on all ice types combined from 1985-1987 to 1996-1999. A Poisson regression model indicated highly significant density declines of 72% on shorefast ice and 43% on pack ice over the 15-year period. However, the apparent decline between the mid-1980s and the late 1990s may have been due to a difference in the timing of surveys rather than an actual decline in abundance (Frost et al. 2002, Kelly et al. 2006). As these surveys represent only a fraction of the stock's range and occurred more than a decade ago, current and reliable data on trends in population abundance for the Alaska stock of ringed seals are considered unavailable.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

A reliable estimate of the maximum net productivity rate is currently unavailable for the Alaska stock of ringed seals. Hence, until additional data become available, it is recommended that the pinniped maximum theoretical net productivity rate (R_{MAX}) of 12% be employed for this stock (Wade and Angliss 1997).

POTENTIAL BIOLOGICAL REMOVAL

Under the 1994 reauthorized Marine Mammal Protection Act (MMPA), the potential biological removal (PBR) is defined as the product of the minimum population estimate, one-half the maximum theoretical net productivity rate, and a recovery factor: $PBR = N_{MIN} \times 0.5R_{MAX} \times F_R$. The recovery factor (F_R) for this stock is 0.5, the value for pinniped stocks with unknown population status (Wade and Angliss 1997). Since the data used to produce the abundance estimate presented in Kelly et al. (2010a) are more than 8 years old, and no reliable N_{MIN} is available, PBR is undetermined.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

New Serious Injury Guidelines

NMFS updated its serious injury designation and reporting process, which uses guidance from previous serious injury workshops, expert opinion, and analysis of historic injury cases to develop new criteria for distinguishing serious from non-serious injury (Angliss and DeMaster 1998, Andersen *et al.* 2008, NOAA 2012). NMFS defines serious injury as an “injury that is more likely than not to result in mortality.” Injury determinations for stock assessments revised in 2013 or later incorporate the new serious injury guidelines, based on the most recent 5-year period for which data are available.

Fisheries Information

Until 2003, there were three different federally-regulated commercial fisheries in Alaska that could have interacted with ringed seals and were monitored for incidental mortality by fishery observers. As of 2003, changes in fishery definitions in the List of Fisheries have resulted in separating these three fisheries into 12 fisheries (69 FR 70094, 2 December 2004). This change does not represent a change in fishing effort, but provides managers with better information on the component of each fishery that is responsible for the incidental serious injury or mortality of marine mammal stocks in Alaska. Between 2007 and 2011, there were incidental serious injuries and mortalities of ringed seals in the Bering Sea/Aleutian Islands flatfish trawl fishery, the Bering Sea/ Aleutian Islands pollock trawl, Bering Sea/ Aleutian Islands Pacific cod trawl, and the Bering Sea/ Aleutian Islands Pacific cod longline (Table 24). Based on data from 2007 to 2011, there have been an average of 3.52 (CV = 0.06) mortalities of ringed seals incidental to commercial fishing operations.

Table 24. Summary of incidental mortality of ringed seals (Alaska stock) due to commercial fisheries from 2007 to 2011 and calculation of the mean annual mortality rate (Breiwick 2013). Details of how percent observer coverage is measured is included in Appendix 6.

Fishery name	Years	Data type	Observer coverage	Observed mortality (in given yrs.)	Estimated mortality (in given yrs.)	Mean annual mortality
Bering Sea/Aleutian Is. flatfish trawl	2007	obs	72	0	0	2.0 (CV = 0.02)
	2008	data	100	2	2.0	
	2009		100	1	1.0	
	2010		100	0	0	
	2011		100	6 (+1)*	6.0 (7)**	
Bering Sea/ Aleutian Is. pollock trawl	2007	obs	85	0	0	1.0 (CV = 0.04)
	2008	data	85	1	1.0	
	2009		86	1	1.0	
	2010		86	0	0	
	2011		98	3	3.0	
Bering Sea/ Aleutian Is. Pacific cod trawl	2007	obs	53	0	0	0.2 (CV = 0.01)
	2008	data	59	0	0	
	2009		63	0	0	
	2010		66	0	0	
	2011		60	1	1.0	

Fishery name	Years	Data type	Observer coverage	Observed mortality (in given yrs.)	Estimated mortality (in given yrs.)	Mean annual mortality
Bering Sea/ Aleutian Is. Pacific cod longline	2007	obs	63	0	0	0.32 (CV = 0.6)
	2008	data	63	0	0	
	2009		60	0	0	
	2010		64	0	0	
	2011		57	1	1.6	
Total estimated annual mortality						3.52 (CV = 0.06)

*Total mortalities observed in unsampled hauls.

**Total mortalities observed in sampled and unsampled hauls. Since the total known mortality (7) exceeds the estimated mortality (6.0) for 2011, the sum of actual mortalities observed (7) will be used as a minimum estimate for that year.

Subsistence/Native Harvest Information

Ringed seals are an important species for Alaska Native subsistence hunters. The estimated annual subsistence harvest in Alaska dropped from 7,000 to 15,000 in the period from 1962 to 1972 to an estimated 2,000 to 3,000 in 1979 (Frost 1985). Based on data from two villages on St. Lawrence Island, the annual take in Alaska during the mid-1980s likely exceeded 3,000 seals (Kelly 1988a).

The Division of Subsistence, Alaska Department of Fish and Game (ADFG), maintained a database that provided additional information on the subsistence harvest of ice seals in different regions of Alaska (ADFG 2000a, b). Information on subsistence harvest of ringed seals was compiled for 129 villages from reports from the Division of Subsistence (Coffing et al. 1998, Georgette et al. 1998, Wolfe and Hutchinson-Scarborough 1999) and a report from the Eskimo Walrus Commission (Sherrod 1982). Data were lacking for 22 villages; their harvests were estimated using the annual per capita rates of subsistence harvest from a nearby village. Harvest levels were estimated from data gathered in the 1980s for 16 villages; otherwise, data gathered from 1990 to 1998 were used. As of August 2000, the subsistence harvest database indicated that the estimated number of ringed seals harvested for subsistence use per year is 9,567. Data on community subsistence harvests are no longer routinely being collected, and no new statewide annual harvest estimates exist. Five Alaska Native communities in the Northwest Arctic region of Alaska voluntarily reported a total of 40 ringed seals were harvested during 2012 (Ice Seal Committee, 2013).

At this time, there are no efforts to quantify the total statewide level of harvest of ringed seals by all Alaska communities. A report on ice seal subsistence harvest in three Alaskan communities indicated that the number and species of ice seals harvested in a particular village may vary considerably between years (Coffing et al. 1999). These interannual differences are likely due to differences in ice and wind conditions that change the hunters' access to different ice habitats frequented by different types of seals. Regardless of the extent to which the harvest may vary interannually, it is clear that the harvest level of 9,567 ringed seals estimated by the Division of Subsistence is considerably higher than the previous minimum estimate reported by Frost (1985). Although some of the more recent entries in the ADFG database have associated measures of uncertainty (Coffing et al. 1999, Georgette et al. 1998), the overall total does not. The estimate of 9,567 ringed seals is the best estimate currently available.

Other human-caused mortality and injury

Between 2007 and 2011, there were 4 records of dead and injured ringed seals reported to the Alaska Regional Office Marine Mammal Stranding Network. All 3 injuries were considered not serious (Allen et al. 2014). One male ringed seal was found in 2008 with a packing band and circumferential wound around its neck; it was disentangled. Two injured ringed seals were reported in 2010, one with a bleeding flipper that was captured and released on site, another that was caught in a subsistence salmon set net. This animal was disentangled by ADFG and released. In 2011, one ringed seal was reported dead from a gunshot wound to the head, presumably a struck and lost animal from the subsistence hunt. This animal presented with skin lesions consistent with those seen in animals considered part of the multi-species northern pinnipeds 2011 Unusual Mortality Event.

STATUS OF STOCK

On December 28, 2012, NMFS listed Arctic ringed seals (*Phoca hispida hispida*) and, thus, the Alaska stock of ringed seals, as threatened under the Endangered Species Act (ESA) (77 FR 76706). The primary concern for this population is the ongoing and anticipated loss of sea ice and snow cover stemming from climate change, which is expected to pose a significant threat to the persistence of these seals in the foreseeable future (based on projections through the end of the 21st century; Kelly et al. 2010a). Because of its threatened status under the ESA, this stock is designated as “depleted” under the MMPA. As a result, the stock is classified as a strategic stock. Since PBR is undetermined, it is not known whether current annual level of incidental U. S. commercial fishery-related mortality (3.5) exceeds 10% of the PBR. However, mortality and serious injury incidental to commercial fishing is likely insignificant and approaching a zero mortality and serious injury rate. The total estimated level of human-caused mortality and serious injury based on commercial fisheries observer data (3.5) and Alaska Native harvest (9,567) is 9,571. Population trends and status of this stock relative to OSP are currently unknown.

Habitat Concerns

The main concern about the conservation status of ringed seals stems from the likelihood that their sea-ice and snow habitats have been modified by the warming climate and, more so, that the scientific consensus projections are for continued and perhaps accelerated warming in the foreseeable future (Kelly et al. 2010a). Climate models consistently project overall diminishing ice and snow cover through the 21st century with regional variation in the timing and severity of those losses. Increasing atmospheric concentrations of greenhouse gases are driving climate warming and increasing acidification of the ringed seal’s habitat. Changes in ocean temperature, acidification, and ice cover threaten prey communities on which ringed seals depend. Laidre et al. (2008) concluded that on a worldwide basis ringed seals were likely to be highly sensitive to climate change based on an analysis of various life history features that could be affected by climate.

The greatest impacts to ringed seals from diminished ice cover will be mediated through diminished snow accumulation. While winter precipitation is forecasted to increase in a warming Arctic (Walsh et al. 2005), the duration of ice cover will be substantially reduced, and the net effect will be lower snow accumulation on the ice (Hezel et al. 2012). Ringed seals excavate subnivean lairs (snow caves) in drifts over their breathing holes in the ice, in which they rest, give birth, and nurse their pups for 5-9 weeks during late winter and spring (Chapskii 1940, McLaren 1958, Smith and Stirling 1975). Snow depths of at least 50-65 cm are required for functional birth lairs (Smith and Stirling 1975, Lydersen and Gjertz 1986, Kelly 1988b, Lydersen 1998, Lukin et al. 2006), and such depths typically are found only where 20-30 cm or more of snow has accumulated on flat ice and then drifted along pressure ridges or ice hummocks (Lydersen et al. 1990, Hammill and Smith 1991, Lydersen and Ryg 1991, Smith and Lydersen 1991). According to climate model projections, snow cover is forecasted to be inadequate for the formation and occupation of birth lairs within this century over the Alaska stock’s entire range (Kelly et al. 2010a). Without the protection of the lairs, ringed seals—especially newborns—are vulnerable to freezing and predation (Kumlien 1879, McLaren 1958, Lukin and Potelov 1978, Smith and Hammill 1980, Lydersen and Smith 1989, Stirling and Smith 2004). Changes in the ringed seal’s habitat will be rapid relative to their generation time and, thereby, will limit adaptive responses. As ringed seal populations decline, the significance of currently lower-level threats—such as ocean acidification, increases in human activities, and changes in populations of predators, prey, competitors, and parasites—may increase.

Additional habitat concerns include the potential effects from oil and gas exploration activities, particularly in the outer continental shelf leasing areas, such as disturbance from vessel traffic, seismic exploration noise, or the potential for oil spills.

CITATIONS

- Alaska Department of Fish and Game. 2000a. Community Profile Database 3.04 for Access 97. Division of Subsistence, Anchorage.
- Alaska Department of Fish and Game. 2000b. Seals+ Database for Access 97. Division of Subsistence, Anchorage.
- Allen, B. M., V. T. Helker, and L. A. Jemison. 2014. Human-caused injury and mortality of NMFS-managed Alaska marine mammal stocks, 2007-2011. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-274, 84 p.
- Bengtson, J. L., L. M. Hiruki-Raring, M. A. Simpkins, and P. L. Boveng. 2005. Ringed and bearded seal densities in the eastern Chukchi Sea, 1999-2000. *Polar Biol.* 28: 833-845.
- Breiwick, J. M. 2013. North Pacific marine mammal bycatch estimation methodology and results, 2007-2011. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-260, 40 p.

- Burns, J. J. 1970. Remarks on the distribution and natural history of pagophilic pinnipeds in the Bering and Chukchi Seas. *J. Mammal.* 51:445-454.
- Burns, J. J., and S. J. Harbo. 1972. An aerial census of ringed seals, northern coast of Alaska. *Arctic* 25:279-290.
- Chapkskii, K. K. 1940. The ringed seal of western seas of the Soviet Arctic (The morphological characteristic, biology and hunting production). Page 147 in N. A. Smirnov, editor. Proceedings of the Arctic Scientific Research Institute, Chief Administration of the Northern Sea Route. Izd. Glavsevmorputi, Leningrad, Moscow. (Translated from Russian by the Fisheries Research Board of Canada, Ottawa, Canada, Translation Series No. 1665, 147 p.).
- Coffing, M., C. Scott, and C.J. Utermohle. 1998. The subsistence harvest of seals and sea lions by Alaska Natives in three communities of the Yukon-Kuskokwim Delta, Alaska, 1997-1998. Technical Paper No. 255, Alaska Dep. Fish and Game, Division of Subsistence, Juneau.
- Coffing, M., C. Scott, and C.J. Utermohle. 1999. The subsistence harvest of seals and sea lions by Alaska Natives in three communities of the Yukon-Kuskokwim Delta, Alaska, 1998-1999. Technical Paper No. 257, Alaska Dep. Fish and Game, Division of Subsistence, Juneau.
- Crawford, J. A., K. J. Frost, L. T. Quakenbush, and A. Whiting. 2012. Different habitat use strategies by subadult and adult ringed seals (*Phoca hispida*) in the Bering and Chukchi seas. *Polar Biology* 35:241-255.
- Freitas, C., K. M. Kovacs, R. A. Ims, M. A. Fedak, and C. Lydersen. 2008. Ringed seal post-moulting movement tactics and habitat selection. *Oecologia* 155:193-204.
- Frost, K. J. 1985. The ringed seal (*Phoca hispida*). Pages 79-87 in J. J. Burns, K. J. Frost, and L. F. Lowry, editors. Marine Mammals Species Accounts. Alaska Department Fish and Game, Juneau, AK.
- Frost, K. J., and L. F. Lowry. 1984. Trophic relationships of vertebrate consumers in the Alaskan Beaufort Sea. Pages 381-401 in P. W. Barnes, D. M. Schell, and E. Reimnitz, editors. The Alaskan Beaufort Sea -- Ecosystems and Environments. Academic Press, Inc., New York, NY.
- Frost, K. J., L. F. Lowry, J. R. Gilbert, and J. J. Burns. 1988. Ringed seal monitoring: relationships of distribution and abundance to habitat attributes and industrial activities. Final Rep. contract no. 84-ABC-00210 submitted to U.S. Dep. Interior, Minerals Management Service, Anchorage, AK. 101 pp.
- Frost, K. J., L. F. Lowry, G. Pendleton, and H. R. Nute. 2002. Monitoring distribution and abundance of ringed seals in northern Alaska. OCS Study MMS 2002-04. Final report from the Alaska Dep. Fish and Game, Juneau, AK, for U.S. Minerals Management Service, Anchorage, AK. 66 pp. + Appendices.
- Frost, K. J., L. F. Lowry, G. Pendleton, and H. R. Nute. 2004. Factors affecting the observed densities of ringed seals, *Phoca hispida*, in the Alaskan Beaufort Sea, 1996-99. *Arctic* 57:115-128.
- Georgette, S., M. Coffing, C. Scott, and C. Utermohle. 1998. The subsistence harvest of seals and sea lions by Alaska Natives in the Norton Sound-Bering Strait Region, Alaska, 1996-97. Technical Paper No. 242, Alaska Dep. Fish and Game, Division of Subsistence, Juneau.
- Hammill, M. O., and T. G. Smith. 1991. The role of predation in the ecology of the ringed seal in Barrow Strait, Northwest Territories, Canada. *Marine Mammal Science* 7:123-135.
- Hezel, P. J., X. Zhang, C. M. Bitz, B. P. Kelly, and F. Massonnet. 2012. Projected decline in spring snow depth on Arctic sea ice caused by progressively later autumn open ocean freeze-up this century. *Geophysical Research Letters* 39:L17505.
- Ice Seal Committee. 2013. Northwest Arctic ice seal harvest survey for 2012. Ice Seal Committee Newsletter, 1(1):3.
- Kelly, B. P. 1988a. Ringed seal, *Phoca hispida*. Pp. 57-75 In J. W. Lentfer (ed.), Selected marine mammals of Alaska. Species accounts with research and management recommendations. Marine Mammal Commission, Washington, D.C.
- Kelly, B. P. 1988b. Locating and characterizing ringed seal lairs and breathing holes in coordination with surveys using forward looking infra-red sensors Fisheries and Oceans Freshwater Institute Final Report. 17 p.
- Kelly, B. P., O. H. Badajos, M. Kunasranta, and J. Moran. 2006. Timing and re-interpretation of ringed seal surveys. Coastal Marine Institute University of Alaska Fairbanks, Final Report. 60 p.
- Kelly, B. P., M. Ponce, D. A. Tallmon, B. J. Swanson, and S. K. Sell. 2009. Genetic diversity of ringed seals sampled at breeding sites; implications for population structure and sensitivity to sea ice loss. University of Alaska Southeast, North Pacific Research Board 631 Final Report. 28 p.
- Kelly, B. P., J. L. Bengtson, P. L. Boveng, M. F. Cameron, S. P. Dahle, J. K. Jansen, E. A. Logerwell, J. E. Overland, C. L. Sabine, G. T. Waring, and J. M. Wilder. 2010a. Status review of the ringed seal (*Phoca hispida*). U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-212, 250 p.

- Kelly, B. P., O. H. Badajos, M. Kunasranta, J. R. Moran, M. Martinez-Bakker, D. Wartzok, and P. Boveng. 2010b. Seasonal home ranges and fidelity to breeding sites among ringed seals. *Polar Biol.* 33:1095-1109.
- King, J. E. 1983. Seals of the world. 2nd ed. Br. Muss. (Nat. Hits.), London. 240 pp.
- Kumlien, L. 1879. Mammals. Pages 55-61 in Contributions to the Natural History of Arctic America made in connection with the Howgate Polar Expedition 1877-78. Government Printing Office, Washington, D.C.
- Laidre, K. L., I. Stirling, L. Lowry, Ø. Wiig, M. P. Heide-Jørgensen, and S. Ferguson. 2008. Quantifying the sensitivity of arctic marine mammals to climate-induced habitat change. *Ecol. Appl.* 18(2):S97-S125.
- Lukin, L. P., G. N. Ogetov, and N. S. Boiko. 2006. Ecology of the ringed seal in the White Sea. UrO RAN, Ekaterinburg, Russia. 165 p. (Translated from Russian by the Baltic Fund for Nature (BFN), State University of St. Petersburg, Russia).
- Lukin, L. R., and V. A. Potelov. 1978. Living conditions and distribution of ringed seal in the White Sea in the winter. *Soviet J. Mar. Biol.* 4:684-690.
- Lydersen, C. 1998. Status and biology of ringed seals (*Phoca hispida*) in Svalbard. Pages 46-62 in M. P. Heide-Jørgensen and C. Lydersen, editors. Ringed Seals in the North Atlantic. NAMMCO Scientific Publications, Volume 1, Tromsø, Norway.
- Lydersen, C., and I. Gjertz. 1986. Studies of the ringed seal (*Phoca hispida* Schreber 1775) in its breeding habitat in Kongsfjorden, Svalbard. *Polar Res.* 4:57-63.
- Lydersen, C., P. M. Jensen, and E. Lydersen. 1990. A survey of the Van Mijen Fiord, Svalbard, as habitat for ringed seals, *Phoca hispida*. *Holarctic Ecol.* 13:130-133.
- Lydersen, C., and M. Ryg. 1991. Evaluating breeding habitat and populations of ringed seals *Phoca hispida* in Svalbard fjords. *Polar Rec.* 27:223-228.
- Lydersen, C., and T. G. Smith. 1989. Avian predation on ringed seal *Phoca hispida* pups. *Polar Biol.* 9:489-490.
- McLaren, I. A. 1958. The biology of the ringed seal (*Phoca hispida* Schreber) in the eastern Canadian Arctic. *Bull. Fish. Res. Bo. Can.* 118:97.
- Moulton, F. D., W. J. Richardson, T. L. McDonald, R. E. Elliott, and M. T. Williams. 2002. Factors influencing local abundance and haulout behavior of ringed seals (*Phoca hispida*) on landfast ice of the Alaskan Beaufort Sea. *Can. J. Zool.* 80:1900-1917.
- NOAA. 2012. Federal Register 77:3233. National Policy for Distinguishing Serious From Non-Serious Injuries of Marine Mammals. <http://www.nmfs.noaa.gov/op/pds/documents/02/238/02-238-01.pdf>.
- Palo, J. 2003. Genetic diversity and phylogeography of landlocked seals. Dissertation. University of Helsinki, Helsinki, Finland. 29 p.
- Palo, J. U., H. S. Mäkinen, E. Helle, O. Stenman, and R. Väinölä. 2001. Microsatellite variation in ringed seals (*Phoca hispida*): genetic structure and history of the Baltic Sea population. *Heredity* 86:609-617.
- Sherrod, G.K. 1982. Eskimo Walrus Commission's 1981 Research Report: The Harvest and Use of Marine Mammals in Fifteen Eskimo Communities. Kawerak, Inc., Nome.
- Smith, T. G., and M. O. Hammill. 1980. A survey of the breeding habitat of ringed seals and a study of their behavior during the spring haul-out period in southeastern Baffin Island. Addendum to the Final Report to the Eastern Arctic Marine Environmental Studies (EAMES) project. Department of Fisheries and Oceans, Arctic Biological Station, Canadian Manuscript Report of Fisheries and Aquatic Sciences, No. 1561. 47p.
- Smith, T. G., and C. Lydersen. 1991. Availability of suitable land-fast ice and predation as factors limiting ringed seal populations, *Phoca hispida*, in Svalbard. *Polar Res.* 10:585-594.
- Smith, T. G., and I. Stirling. 1975. The breeding habitat of the ringed seal (*Phoca hispida*). The birth lair and associated structures. *Can. J. Zool.* 53:1297-1305.
- Stirling, I., and T. G. Smith. 2004. Implications of warm temperatures, and an unusual rain event for the survival of ringed seals on the coast of southeastern Baffin Island. *Arctic* 57:59-67.
- Wade, P. R., and R. Angliss. 1997. Guidelines for assessing marine mammal stocks: report of the GAMMS workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Walsh, J. E., O. Anisimov, J. O. M. Hagen, T. Jakobsson, J. Oerlemans, T. D. Prowse, V. Romanovsky, N. Savelieva, M. Serreze, A. Shiklomanov, I. Shiklomanov, and S. Solomon. 2005. Section 6.2. Precipitation and evapotranspiration. Pages 184-189 in Arctic Climate Impact Assessment. Cambridge University Press, Cambridge, UK.
- Wolfe, R., and L.B. Hutchinson-Scarborough. 1999. The subsistence harvest of harbor seal and sea lion by Alaska Natives in 1998. Technical Paper No. 250, Alaska Dep. Fish and Game, Division of Subsistence, Juneau.