

SHSTMP_PS_Delta_Habitat_Features_2011.shp data dictionary

This layer was developed by NOAA Fisheries to delineate distributaries, tidal channels, tidal channel complexes, and tidal mud flats of major deltas within the Puget Sound region to be used as part of salmon and steelhead habitat monitoring effort by Salmon Habitat Status and Trend Monitoring Program (SHSTMP). Habitat features were digitized at 1:1000 scale using 0.3-meter resolution true-color Microsoft Global Ortho project aerial imagery with positional accuracy of 4 meters. Imagery was collected from July 8, 2010 to August 1, 2011 using Ultracam-G digital camera on an aircraft flown at an altitude of 5,000 m.

Channel was identified as a distributary when; 1) its bankfull width was greater than 2-3 meters, 2) there was a clear upstream connection to the main channel or other distributaries, and 3) the channel had a partial or complete continuous flow path from the river to the delta mouth.

Distributaries were designated as primary when the size of channels at bifurcations indicated the greatest amount of river flow to be moving downstream. We digitized the perimeter of all distributaries along the bankfull edge

Channel was identified as tidal channel when there was no clear connection at the upstream end. Tidal channels were designated as large where bankfull width was at least 2-3 meters, and small where bankfull width was less than 2-3 meters. We digitized the perimeter of all large tidal channels along the vegetated channel edge or the artificial bank edge (along docks, seawalls, etc.) where connection between distributaries, tidal channels, or the geomorphic delta boundary was evident from the aerial image. For small tidal channels, we digitized polylines along the flow path and then buffered the polylines by 1 meter to create a polygon feature.

Where mostly vegetated marsh and scrub shrub environments prevent accurate delineation of tidal channels, we digitized polygons around tidal channel complexes. Polygons were digitized to include both the vegetated islands and tidal channel network. Tidal complexes were also digitized in areas of maturing restoration projects, where vegetation has become mostly established, but channels have not yet fully formed.

Tidal flat perimeter was digitized within the delta polygons where complex channel networks occurred within largely unvegetated, tidally flooded areas. We restricted the delineation of tidal flats to the seaward extent of vegetated marsh and excluded mud flat habitat that occurred at the delta terminus. Interior delta areas within restoration project extents were also digitized as tidal flats where tidal connectivity was restored but channel formation and vegetation establishment have not progressed to develop clearly defined channel networks between vegetated substrate.

Where it was evident that a habitat feature was created as part of a restoration project, it was designated as restoration. Similarly, where it was evident that a habitat feature was constructed or heavily modified (often with piers or docks), it was designated as modified.

Field Name	Description	Units
Delta	Delta name: DOS – Dosewallips DUC – Duckabush DUN – Dungeness DUW – Duwamish ELW – Elwha HAM – Hamma Hamma NKS – Nooksack NSQ – Nisqually PUY – Puyallup QUL – Big Quilcene SAM – Sammamish SKG – Skagit SKO – Skokomish SNO - Snohomish STL – Stillaguamish SWI – Swinomish UNI – Union	
F_Type	Feature type: Distributary, Tidal Channel, Tidal Complex, Tidal Flat	
S_Type	Channel size designation. Distributaries: Primary, Secondary Tidal channels: Large, Small	
M_Type	Feature modification designation: Modified, Restoration.	
Image_Date	Aerial imagery collection date	
Ck_MPG	Puget Sound Chinook salmon major population groups (NMFS, 2007): Central/South Basin Hood Canal Strait of Georgia Strait of Juan de Fuca Whidbey Basin	
Stl_MPG	Puget Sound steelhead salmon major population groups (NMFS, 2011): Northern Cascades Olympic South-Central Cascades	
Area_ha	Polygon area	hectare
Perim_km	Perimeter length	kilometer

References

- NMFS (National Marine Fisheries Service). 2007. Puget Sound Salmon Recovery Plan, volume 1. Shared Strategy for Puget Sound, Seattle.
- NMFS (National Marine Fisheries Service). 2011. 5-Year Review: Summary & Evaluation of Puget Sound Chinook, Hood Canal Summer Chum, Puget Sound Steelhead. NMFS Northwest Region, Portland, Oregon.