


## BLUE WHALE (*Balaenoptera musculus musculus*): Central North Pacific Stock

### STOCK DEFINITION AND GEOGRAPHIC RANGE

The International Whaling Commission (IWC) has formally considered only one management stock for blue whales in the North Pacific (Donovan 1991), but up to five populations have been proposed (Reeves et al. 1998). Rice (1974) hypothesized that blue whales from Baja California migrated far offshore to feed in the eastern Aleutians or Gulf of Alaska and returned to feed in California waters; however, he has more recently concluded that the California population is separate from the Gulf of Alaska population (Rice 1992). Length frequency analyses (Gilpatrick et al. 1996) and photo-identification studies (Calambokidis et al. 1995) support separate population status for blue whales feeding off California and those feeding in Alaskan waters. Whaling catch data indicate that whales feeding along the Aleutian Islands are probably part of a central Pacific stock (Reeves et al. 1998), which may migrate to offshore waters north of Hawaii in winter (Berzin and Rovnin 1966). Blue whale feeding aggregations have not been found in Alaska despite several surveys (Leatherwood et al. 1982; Stewart et al. 1987; Forney and Brownell 1996); however, blue whale calls have been recorded there between 1995 and 2001 (Stafford et al. 2001, Stafford 2003).

Recent analyses of acoustic data obtained throughout the North Pacific Ocean (Stafford et al. 2001; Stafford 2003) has revealed two distinct blue whale call types, suggesting two North Pacific stocks: eastern and central (formerly western). The regional occurrence patterns indicate that blue whales from the eastern North Pacific stock winter off Mexico, central America, and as far south as 8° S (Stafford et al. 1999), and feed during summer off the U. S. West Coast and to a lesser extent in the Gulf of Alaska, . This stock has previously been documented to feed in waters off California (and occasionally as far north as British Columbia; Calambokidis et al. 1998) in summer/fall (from June to November) migrating south to productive areas off Mexico (Calambokidis et al. 1990) and as far south as the Costa Rica Dome (10° N) in winter/spring (Mate et al. 1999, Stafford et al. 1999). Blue whales belonging to the central Pacific stock appear to feed in summer southwest of Kamchatka, south of the Aleutians, and in the Gulf of Alaska (Stafford 2003; Watkins et al. 2000), and in winter they migrate to lower latitudes in the western Pacific and less frequently in the central Pacific, including Hawaii (Stafford et al. 2001). The only published sighting record of blue whales near Hawaii is that of Berzin and Rovnin (1966). Two sightings have been made by observers on Hawaii-based longline vessels (Figure 1; NMFS/PIR, unpublished data). Additional evidence that blue whales occur in this area comes from acoustic recordings made off Oahu and Midway Islands (Northrop et al. 1971; Thompson and Friedl 1982), which included at least some within the U.S. Exclusive Economic Zone (EEZ). The recordings made off Hawaii showed bimodal peaks throughout the year (Stafford et al. 2001), with central Pacific call types heard during winter and eastern Pacific calls heard during summer. For the Marine Mammal Protection Act (MMPA) stock assessment reports, there are two blue whale stocks within the



**Figure 1.** Locations of four blue whale sightings made by observers aboard Hawaii-based longline fishing vessels between July 1994 and December 2009 (NMFS/PIR unpublished data). Solid lines represent the U.S. Exclusive Economic Zone (EEZ).

Pacific U.S. EEZ: 1) the central North Pacific stock (this report), which includes whales found around the Hawaiian Islands during winter and 2) the eastern North Pacific stock, which feeds primarily off California.

### **POPULATION SIZE**

From ship line-transect surveys, Wade and Gerrodette (1993) estimated 1,400 blue whales for the eastern tropical Pacific. No blue whale sightings were made during a summer 1994 shipboard survey south of the Aleutian Islands (Forney and Brownell 1996), during twelve aerial surveys conducted in 1993-98 within about 25 nmi of the main Hawaiian Islands (Mobley et al. 2000), or during a summer/fall 2002 shipboard surveys of the entire Hawaiian Islands EEZ (Barlow 2006). Therefore, no estimate of abundance is available for the central Pacific blue whale stock.

#### **Minimum Population Estimate**

No data are available to provide a minimum population estimate.

#### **Current Population Trend**

No data are available on current population trend.

### **CURRENT AND MAXIMUM NET PRODUCTIVITY RATES**

No data are available on current or maximum net productivity rate.

### **POTENTIAL BIOLOGICAL REMOVAL**

No PBR can be calculated for this stock at this time.

### **HUMAN-CAUSED MORTALITY AND SERIOUS INJURY**

#### **Fishery Information**

Information on fishery-related mortality of cetaceans in Hawaiian waters is limited, but the gear types used in Hawaiian fisheries are responsible for marine mammal mortality and serious injury in other fisheries throughout U.S. waters. Gillnets appear to capture marine mammals wherever they are used, and float lines from lobster traps and longlines can be expected to occasionally entangle cetaceans (Perrin et al. 1994). Interactions with cetaceans are reported for all pelagic fisheries, and large whales have been entangled in longline gear off the Hawaiian Islands (Nitta and Henderson 1993, Forney 2009). There are currently two distinct longline fisheries based in Hawaii: a deep-set longline (DSL) fishery that targets primarily tunas, and a shallow-set longline fishery (SSL) that targets swordfish. Both fisheries operate within U.S. waters and on the high seas. Between 2004 and 2008, no blue whales were observed hooked or entangled in the SSL fishery (100% observer coverage) or the DSL fishery (20-28% observer coverage) (McCracken & Forney 2010).

#### **Historical Mortality**

At least 9,500 blue whales were taken by commercial whalers throughout the North Pacific between 1910 and 1965 (Ohsumi and Wada 1972). Some proportion of this total may have been from a population or populations that migrate seasonally into the Hawaiian EEZ. The species has been protected in the North Pacific by the IWC since 1966.

### **STATUS OF STOCK**

The status of blue whales in Hawaiian waters relative to OSP is unknown, and there are insufficient data to evaluate trends in abundance. Blue whales are formally listed as "endangered" under the Endangered Species Act (ESA), and consequently the central Pacific stock is automatically considered as a "depleted" and "strategic" stock under the MMPA. Because there have been no reported fishery related mortality or serious injuries of blue whales within the Hawaiian Islands EEZ, the total fishery-related mortality and serious injury of this stock can be considered to be insignificant and approaching zero. Increasing levels of anthropogenic noise in the world's oceans has been suggested to be a habitat concern for blue whales (Reeves et al. 1998).

### **REFERENCES**

- Barlow, J. 2006. Cetacean abundance in Hawaiian waters estimated from a summer/fall survey in 2002. *Marine Mammal Science* 22: 446–464.
- Berzin, A. A. and A. A. Rovnin. 1966. Distribution and migration of whales in the northeastern part of the Pacific Ocean, Bering and Chukchi Seas. *Izv. Tikhookean. Nauchno-issled. Inst. Rybn. Khoz. Okeanogr.*

- (TINRO) 58:179-207. {In Russian}. (Translated by U.S. Dep. Inter., Bur. Commer. Fish., Seattle, Washington, 1966, pp. 103-106 *In*: Panin, K. I. (ed) Soviet Research on marine mammals of the Far East).
- Calambokidis, J, T. Chandler, K. Rasmussen, G.H. Steiger, and L. Schlender. 1998. Humpback and blue whale photographic identification: Report of research in 1997. Final report to Southwest Fisheries Science Center, Olympic Coast National Marine Sanctuary, University of California at Santa Cruz, and Cornell University. Cascadia Research, 218½ W. Fourth Ave., Olympia, WA 98501. 41pp.
- Calambokidis, J., R. Sears, G. H. Steiger, and J. Evenson. 1995. Movement and stock structure of blue whales in the eastern North Pacific. P.19 *In*: Proceedings of the Eleventh Biennial Conference on the Biology of Marine Mammals, Orlando, FL, 14-18 December 1995 (Abstract). Society for Marine Mammalogy, Lawrence, KS.
- Calambokidis, J., G. H. Steiger, J.C. Cabbage, K.C. Balcomb, C. Ewald, S. Kruse, R. Wells, and R. Sears. 1990. Sightings and movement of blue whales off central California 1986-88 from photo-identification of individuals. *Rep. Int. Whaling. Commn., Special Issue 12*:343-348.
- Donovan, G. P. 1991. A review of IWC stock boundaries. *Rept. Int. Whal. Commn., Special Issue 13*:39-68.
- Forney, K.A. 2009. Serious injury determinations for cetaceans caught in Hawaii longline fisheries during 1994-2008. Draft document PSRG-2009-09 presented to the Pacific Scientific Review Group, November 3-5, 2009, Del Mar, CA..
- Forney, K. A. and Brownell, R. L., Jr. 1996. Preliminary report of the 1994 Aleutian Island Marine Mammal Survey. Paper SC/48/O11 presented to the International Whaling Commission, June 1996 (unpublished).
- Gilpatrick, J., W. Perryman, M. Lynn, and M. A. DeAngelis. 1996. Geographic populations of blue whales (*Balaenoptera musculus*) in the North Pacific Ocean investigated from whaling records and aerial photogrammetry. Paper SC/47/NP4 presented to the International Whaling Commission, May 1995 (unpublished).
- Leatherwood, S., R. R. Reeves, W. F. Perrin, and W. E. Evans. 1982. Whales, dolphins, and porpoises of the eastern North Pacific and adjacent Arctic waters: A guide to their identification. NOAA Technical Rept. NMFS Circular 444. 245 pp.
- Mate, B.R., B.A. Lagerquist, and J. Calambokidis. Movements of North Pacific blue whales during the feeding season off southern California and their southern fall migration. *Mar. Mamm. Sci.* 15(4):1246-1257.
- McCracken M., and K.A. Forney. 2010. Preliminary assessment of incidental interactions with marine mammals in the Hawaii longline deep and shallow set fisheries. NMFS, Pacific Islands Fisheries Science Center Working Paper WP-10-001. 27p
- Mobley, J. R. , Jr, S. S. Spitz, K. A. Forney, R. A. Grotendorf, and P. H. Forestall. 2000. Distribution and abundance of odontocete species in Hawaiian waters: preliminary results of 1993-98 aerial surveys *Admin. Rep. LJ-00-14C*. Southwest Fisheries Science Center, National Marine Fisheries Service, P.O. Box 271, La Jolla, CA 92038. 26 pp.
- Nitta, E. and J. R. Henderson. 1993. A review of interactions between Hawaii's fisheries and protected species. *Mar. Fish. Rev.* 55(2):83-92.
- Northrop, J., W. C. Cummings, and M. F. Morrison. 1971. Underwater 20-Hz signals recorded near Midway Island. *J. Acoust. Soc. Am.* 49:1909-1910.
- Ohsumi, S. and S. Wada. 1972. Stock assessment of blue whales in the North Pacific. *Int. Whal. Commn. Sci. Comm. Rep.*, 20 pp.
- Perrin, W.F., G.P. Donovan and J. Barlow. 1994. Gillnets and Cetaceans. *Rep. Int. Whal. Commn., Spec. Iss.* 15, 629pp.
- Reeves, R. R., P. J. Clapham, R. L. Brownell, Jr., and G. K. Silber. 1998. Recovery plan for the blue whale (*Balaenoptera musculus*). Office of Protected Resources, NMFS, NOAA, Silver Spring, Maryland. 30 pp.
- Rice, D. W. 1974. Whales and whale research in the eastern North Pacific. pp. 170-195 *In*: W. E. Schevill (ed.). The Whale Problem: A Status Report. Harvard Press, Cambridge, MA.
- Rice, D. W. 1992. The blue whales of the southeastern North Pacific Ocean. pp. 1-3 *In*. Alaska Fisheries Science Center, *Quart. Rept.* Oct.-Dec.
- Stafford, K.M., S.L. Niekirk, and C.G. Fox. 1999. An acoustic link between blue whales in the eastern tropical Pacific and the northeast Pacific. *Mar. Mamm. Sci.* 15(4):1258-1268.
- Stafford, K.M., S. L. Niekirk, and G.G. Fox. 2001. Geographic and seasonal variation of blue whale calls in the North Pacific. *J. Cetacean Res. Manage.* 3(1):65-76.
- Stafford. K.M. 2003. Two types of blue whale calls recorded in the Gulf of Alaska. *Mar. Mamm. Sci.* 19:682-693
- Stewart, B. S., S. A. Karl, P. K. Yochem, S. Leatherwood, and J. L. Laake. 1987. Aerial surveys for cetaceans in the former Akutan, Alaska, whaling grounds. *Arctic* 40(1):33-42.

- Thompson, P. O. and W. A. Friedl. 1982. A long term study of low frequency sound from several species of whales off Oahu, Hawaii. *Cetology* 45:1-19.
- Wade, P. R. and T. Gerrodette. 1993. Estimates of cetacean abundance and distribution in the eastern tropical Pacific. *Rep. Int. Whal. Commn.* 43:477-493.
- Watkins, W. A., J. E. George, M. A. Daher, K. Mullin, D. L. Martin, S. H. Haga, and N. A. DiMarzio. 2000. Whale call data for the North Pacific November 1995 through July 1999: Occurrence of calling whales and source locations from SOSUS and other acoustic systems. Technical Report WHOI\_00\_02 available from Woods Hole Oceanographic Institution. 160pp.