

HUMPBACK WHALE (*Megaptera novaeangliae*): Gulf of Maine Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

In the western North Atlantic, humpback whales feed during spring, summer and fall over a geographic range encompassing the eastern coast of the United States (including the Gulf of Maine), the Gulf of St. Lawrence, Newfoundland/Labrador, and western Greenland (Katona and Beard 1990). Other North Atlantic feeding grounds occur off Iceland and northern Norway, including off Bear Island, Jan Mayen, and Franz Josef Land (Christensen *et al.* 1992; Palsbøll *et al.* 1997; M. Moore, WHOI, pers. comm.). These six regions represent relatively discrete subpopulations, fidelity to which is determined matrilineally (Clapham and Mayo 1987), which is supported by studies of the mitochondrial genome (Palsbøll *et al.* 1995; Palsbøll *et al.* 2001) and individual animal movements (Stevick *et al.* 2006). In early stock assessment reports, the North Atlantic humpback whale population was treated as a single stock for management purposes (Waring *et al.* 1999). Subsequently, a decision was made to reclassify the Gulf of Maine as a separate feeding stock (Waring *et al.* 2000) based upon the strong fidelity by individual whales to this region, and the attendant assumption that, were this subpopulation wiped out, repopulation by immigration from adjacent areas would not occur on any reasonable management timescale. During the 2002 Comprehensive Assessment of North Atlantic humpback whales, the International Whaling Commission acknowledged the evidence for treating the Gulf of Maine as a separate management unit (IWC 2002).

During the summers of 1998 and 1999, the Northeast Fisheries Science Center conducted surveys for humpback whales on the Scotian Shelf to establish the occurrence and population identity of the animals found in this region, which lies between the well-studied populations of the Gulf of Maine and Newfoundland. Photographs from both surveys were compared to both the overall North Atlantic Humpback Whale Catalogue and a large regional catalogue from the Gulf of Maine (maintained by the College of the Atlantic and the Center for Coastal Studies, respectively); this work is summarized in Clapham *et al.* (2003). The match rate between the Scotian Shelf and the Gulf of Maine was 27% (14 of 52 Scotian Shelf individuals from both years). Comparable rates of exchange were obtained from the southern (28%, $n=10$ of 36 whales) and northern (27%, $n=4$ of 15 whales) ends of the Scotian Shelf, despite the additional distance of nearly 100 nautical miles (one whale was observed in both areas). In contrast, all of the 36 humpback whales identified by the same NMFS surveys elsewhere in the Gulf of Maine (including Georges Bank, southwestern Nova Scotia and the Bay of Fundy) had been previously observed in the Gulf of Maine region. The sighting histories of the 14 Scotian Shelf whales matched to the Gulf of Maine suggested that many of them were transient through the latter area. There were no matches between the Scotian Shelf and any




Figure 1. Distribution of humpback whale sightings from NEFSC and SEFSC shipboard and aerial surveys during the summers of 1995, 1998, 1999, 2002, 2004, 2006, 2007, 2008, 2010 and 2011. Isobaths are the 100-m, 1000-m and 4000-m depth contours.

other North Atlantic feeding ground, except the Gulf of Maine; however, instructive comparisons are compromised by the often low sampling effort in other regions in recent years. Overall, it appears that the northern range of many members of the Gulf of Maine stock does not extend onto the Scotian Shelf.

During winter, whales from most North Atlantic feeding areas (including the Gulf of Maine) mate and calve in the West Indies, where spatial and genetic mixing among feeding groups occurs (Katona and Beard 1990; Clapham *et al.* 1993; Palsbøll *et al.* 1997; Stevick *et al.* 1998). A few whales likely using eastern North Atlantic feeding areas migrate to the Cape Verde Islands (Reiner *et al.* 1996; Wenzel *et al.* 2009). In the West Indies, the majority of whales are found in the waters of the Dominican Republic, notably on Silver Bank and Navidad Bank, and in Samana Bay (Balcomb and Nichols 1982; Whitehead and Moore 1982; Mattila *et al.* 1989, 1994). Humpback whales are also found at much lower densities throughout the remainder of the Antillean arc, from Puerto Rico to the coast of Venezuela (Winn *et al.* 1975; Levenson and Leapley 1978; Price 1985; Mattila and Clapham 1989). Although recognition of 2 breeding areas for North Atlantic humpbacks is the prevailing model, several observations suggest that our knowledge of breeding season distribution is far from complete (see Smith and Pike 2009).

All whales from this stock may not migrate to the West Indies every winter, because significant numbers of animals may be found in mid- and high-latitude regions at this time (Clapham *et al.* 1993; Swingle *et al.* 1993) and some individuals have been resighted across a winter season (Clapham *et al.* 1993; Robbins 2007). Acoustic recordings made in Stellwagen Bank National Marine Sanctuary in 2006 and 2008 detected humpback song in almost all months, including throughout the winter (Vu *et al.* 2012). This confirms the presence of male humpback whales in the area (a mid-latitude feeding ground) through the winter in these years. In addition, photographic records from Newfoundland have shown a number of adult humpbacks remain there year-round, particularly on the island's north coast. In collaboration with colleagues in the French islands of St. Pierre and Miquelon, a new photographic catalogue and concurrent matching effort is being undertaken for this region (J. Lawson, DFO, pers. comm.).

An increased number of sightings of humpback whales in the vicinity of the Chesapeake and Delaware Bays occurred in 1992 (Swingle *et al.* 1993). Wiley *et al.* (1995) reported that 38 humpback whale strandings occurred during 1985–1992 in the U.S. mid-Atlantic and southeastern states. Humpback whale strandings increased, particularly along the Virginia and North Carolina coasts, and most stranded animals were sexually immature; in addition, the small size of many of these whales strongly suggested that they had only recently separated from their mothers. Wiley *et al.* (1995) concluded that these areas were becoming an increasingly important habitat for juvenile humpback whales and that anthropogenic factors may negatively impact whales in this area. There have also been a number of wintertime humpback sightings in coastal waters of the southeastern U.S. Whether the increased numbers of sightings represent a distributional change, or are simply due to an increase in sighting effort and/or whale abundance, is unknown.

A key question with regard to humpback whales off the southeastern and mid-Atlantic states is their population identity. This topic was investigated using fluke photographs of living and dead whales observed in the region (Barco *et al.* 2002). In this study, photographs of 40 whales (alive or dead) were of sufficient quality to be compared to catalogs from the Gulf of Maine (i.e., the closest feeding ground) and other areas in the North Atlantic. Of 21 live whales, 9 (43%) matched to the Gulf of Maine, 4 (19%) to Newfoundland, and 1 (4.8%) to the Gulf of St Lawrence. Of 19 dead humpbacks, 6 (31.6%) were known Gulf of Maine whales. Although the population composition of the mid-Atlantic is apparently dominated by Gulf of Maine whales, lack of photographic effort in Newfoundland makes it likely that the observed match rates under-represent the true presence of Canadian whales in the region. A new photographic catalog and concurrent matching effort is being undertaken for this region which may improve knowledge in this regard. Barco *et al.* (2002) suggested that the mid-Atlantic region primarily represents a supplemental winter feeding ground used by humpbacks.

In New England waters, feeding is the principal activity of humpback whales, and their distribution in this region has been largely correlated to abundance of prey species, although behavior and bathymetry are factors influencing foraging strategy (Payne *et al.* 1986, 1990). Humpback whales are frequently piscivorous when in New England waters, feeding on herring (*Clupea harengus*), sand lance (*Ammodytes* spp.), and other small fishes. In the northern Gulf of Maine, euphausiids are also frequently taken (Paquet *et al.* 1997). Commercial depletion of herring and mackerel led to an increase in sand lance in the southwestern Gulf of Maine in the mid-1970s, with a concurrent decrease in humpback whale abundance in the northern Gulf of Maine. Humpback whales were densest over the sandy shoals in the southwestern Gulf of Maine favored by the sand lance during much of the late 1970s and early 1980s, and humpback distribution appeared to have shifted to this area (Payne *et al.* 1986). An apparent reversal began in the mid-1980s, and herring and mackerel increased as sand lance again decreased (Fogarty *et al.* 1991). Humpback whale abundance in the northern Gulf of Maine increased markedly during 1992–1993, along with a

major influx of herring (P. Stevick, pers. comm.). Humpback whales were few in nearshore Massachusetts waters in the 1992–1993 summer seasons. They were more abundant in the offshore waters of Cultivator Shoal and on the Northeast Peak on Georges Bank and on Jeffreys Ledge; these latter areas are traditional locations of herring occurrence. In 1996 and 1997, sand lance and therefore humpback whales were once again abundant in the Stellwagen Bank area. However, unlike previous cycles, when an increase in sand lance corresponded to a decrease in herring, herring remained relatively abundant in the northern Gulf of Maine, and humpbacks correspondingly continued to occupy this portion of the habitat, where they also fed on euphausiids (Wienrich *et al.* 1997). Diel patterns in humpback foraging behavior have been shown to correlate with diel patterns in sand lance behavior (Friedlaender *et al.* 2009).

In early 1992, a major research program known as the Years of the North Atlantic Humpback (YONAH) (Smith *et al.* 1999) was initiated. This was a large-scale, intensive study of humpback whales throughout almost their entire North Atlantic range, from the West Indies to the Arctic. During two primary years of field work, photographs for individual identification and biopsy samples for genetic analysis were collected from summer feeding areas and from the breeding grounds in the West Indies. Additional samples were collected from certain areas in other years. Results pertaining to the estimation of abundance and to genetic population structure are summarized below.

POPULATION SIZE

North Atlantic Population

The overall North Atlantic population (including the Gulf of Maine), derived from genetic tagging data collected by the YONAH project on the breeding grounds, was estimated to be 4,894 males (95% CI=3,374-7,123) and 2,804 females (95% CI=1,776-4,463) (Palsbøll *et al.* 1997). Because the sex ratio in this population is known to be even (Palsbøll *et al.* 1997), the excess of males is presumed a result of sampling bias, lower rates of migration among females, or sex-specific habitat partitioning in the West Indies; whatever the reason, the combined total is an underestimate of overall population size. Photographic mark-recapture analyses from the YONAH project provided an ocean-basin-wide estimate of 11,570 animals during 1992/1993 (CV=0.068, Stevick *et al.* 2003), and an additional genotype-based analysis yielded a similar but less precise estimate of 10,400 whales (CV=0.138, 95% CI=8,000 to 13,600) (Smith *et al.* 1999).

Gulf of Maine stock - earlier estimates

Please see Appendix IV for earlier estimates. As recommended in the GAMMS Workshop Report (Wade and Angliss 1997), estimates older than eight years are deemed unreliable and should not be used for PBR determinations.

Gulf of Maine Stock - Recent surveys and abundance estimates

An abundance of 335 (CV=0.42) humpback whales was estimated from a line-transect survey conducted during June-August 2011 by ship and plane (Palka 2012). The aerial portion that contributed to the abundance estimate covered 5,313 km of tracklines that were over waters north of New Jersey and shallower than the 100-m depth contour through the U.S. and Canadian Gulf of Maine and up to and including the lower Bay of Fundy. The shipboard portion covered 3,107 km of tracklines that were in waters that were deeper than the 100-m depth contour out to beyond the U.S. EEZ. Both sighting platforms used a two-simultaneous-team data collection procedure, which allows estimation of abundance corrected for perception bias (Laake and Borchers, 2004). Estimation of abundance was based on the independent observer approach assuming point independence (Laake and Borchers 2004) and calculated using the mark-recapture distance sampling option in the computer program Distance (version 6.0, release 2, Thomas *et al.* 2009). This estimate did not include the portion of the Scotian Shelf that is known to be part of the range used by Gulf of Maine humpback whales. These various line-transect surveys lack consistency in geographic coverage, and because of the mobility of humpback whales, pooling stratum estimates across years to produce a single estimate is not advisable. However, similar to an estimate that appeared in Clapham *et al.* (2003), J. Robbins (Center for Coastal Studies, pers. comm.) used photo-id evidence of presence (see Robbins 2009, 2010, 2011 for data description) to calculate the minimum number alive of catalogued individuals seen during the 2008 feeding season within the Gulf of Maine, or seen both before and after 2008, plus whales seen for the first time as non-calves in 2009. That procedure placed the minimum number alive in 2008 at 823 animals.

Minimum Population Estimate

For statistically-based estimates, the minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile

of the log-normal distribution as specified by Wade and Angliss (1997). The most recent line-transect survey, which did not include the Scotian Shelf portion of the stock, produced an estimate of abundance for Gulf of Maine humpback whales of 331 animals (CV=0.48) with a resultant minimum population estimate for this stock of 228 animals. The line-transect based N_{min} is unrealistic because at least 500 uniquely identifiable individual whales from the GOM stock were seen during the calendar year of that survey and the actual population would have been larger because re-sighting rates of GOM humpbacks have historically been <1 (Robbins 2007). Using the minimum count from at least 2 years prior to the year of a stock assessment report allows time to resight whales known to be alive prior to and after the focal year. Thus, the minimum population estimate is set to the 2008 mark-recapture based count of 823.

Table 1. Summary of abundance estimates for Gulf of Maine humpback whales with month, year, and area covered during each abundance survey, and resulting abundance estimate (N_{best}) and coefficient of variation (CV). Note that the second row represents the results from an analysis of resights of individually identified animals.			
Month/Year	Type	N_{best}	CV
Jun-Oct 2008	Gulf of Maine and Bay of Fundy	823	0
Jun-Aug 2011	Virginia to lower Bay of Fundy	335	0.42

Current Population Trend

As detailed below, the most recent available data suggest that the Gulf of Maine humpback whale stock is characterized by a positive trend in size. This is consistent with an estimated average trend of 3.1% (SE=0.005) in the North Atlantic population overall for the period 1979-1993 (Stevick *et al.* 2003), although there are no feeding-area-specific estimates. The best available estimate of the average rate of increase for the West Indies breeding population [which includes the Gulf of Maine feeding stock] is 3.1% per year (SE= 0.005) for the period 1979-1993 (Stevick *et al.* 2003). An analysis of demographic parameters for the Gulf of Maine (Clapham *et al.* 2003) suggested a lower rate of increase than the 6.5% reported by Barlow and Clapham (1997), but results may have been confounded by distribution shifts.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Zerbini *et al.* (2010) reviewed various estimates of maximum productivity rates for humpback whale populations, and, based on simulation studies, they proposed that 11.8% be considered as the maximum rate at which the species could grow. Barlow and Clapham (1997), applying an interbirth interval model to photographic mark-recapture data, estimated the population growth rate of the Gulf of Maine humpback whale stock at 6.5% (CV=0.012). Maximum net productivity is unknown for this population, although a theoretical maximum for any humpback population can be calculated using known values for biological parameters (Brandão *et al.* 2000; Clapham *et al.* 2001). For the Gulf of Maine stock, data supplied by Barlow and Clapham (1997) and Clapham *et al.* (1995) give values of 0.96 for survival rate, 6 years as mean age at first parturition, 0.5 as the proportion of females, and 0.42 for annual pregnancy rate. From this, a maximum population growth rate of 0.072 is obtained according to the method described by Brandão *et al.* (2000). This suggests that the observed rate of 6.5% (Barlow and Clapham 1997) is close to the maximum for this stock.

Clapham *et al.* (2003) updated the Barlow and Clapham (1997) analysis using data from the period 1992 to 2000. The population growth estimate was either 0% (for a calf survival rate of 0.51) or 4.0% (for a calf survival rate of 0.875). Although uncertainty was not strictly characterized by Clapham *et al.* (2003), their work might reflect a decline in population growth rates from the earlier study period.. More recent work by Robbins (2007) places apparent survival of calves at 0.664 (95% CI: 0.517-0.784), a value between those used by Barlow and Clapham (1997) and in addition found productivity to be highly variable and well less than maximum.

Despite the uncertainty accompanying the more recent estimates of observed population growth rate for the Gulf of Maine stock, the maximum net productivity rate was assumed to be 6.5% calculated by Barlow and Clapham (1997) because it represents an observation greater than the default of 0.04 for cetaceans (Barlow *et al.* 1995) but is conservative in that it is well below the results of Zerbini *et al.* (2010).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a recovery factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size for the Gulf of Maine stock is 823 whales. The maximum productivity rate is 0.065. The recovery factor is assumed to be 0.10 because this stock is listed as an endangered species under the Endangered Species Act. PBR for the Gulf of Maine humpback whale stock is 2.7 whales.

ANNUAL HUMAN-CAUSED SERIOUS INJURY AND MORTALITY

For the period 2009 through 2013, the minimum annual rate of human-caused mortality and serious injury to the Gulf of Maine humpback whale stock averaged 9 animals per year. This value includes incidental fishery interaction records, 7.4; and records of vessel collisions, 1.6 (Table 2; Henry *et al.* 2015).

In contrast to stock assessment reports before 2007, these averages include humpback mortalities and serious injuries that occurred in the southeastern and mid-Atlantic states that could not be confirmed as involving members of the Gulf of Maine stock. In past reports, only events involving whales confirmed to be members of the Gulf of Maine stock were counted against the PBR. Starting in the 2007 report, we assumed whales were from the Gulf of Maine unless they were identified as members of another stock. At the time of this writing, no whale was identified as a member of another stock. These determinations may change with the availability of new information. Canadian records from the southern side of Nova Scotia were incorporated into the mortality and serious injury rates, to reflect the effective range of this stock as described above. For the purposes of this report, discussion is primarily limited to those records considered to be confirmed human-caused mortalities or serious injuries.

To better assess human impacts (both vessel collision and gear entanglement) there needs to be greater emphasis on the timely recovery of carcasses and complete necropsies. The literature and review of records described here suggest that there are significant human impacts beyond those recorded in the data assessed for serious injury and mortality. For example, a study of entanglement-related scarring on the caudal peduncle of 134 individual humpback whales in the Gulf of Maine suggested that between 48% and 65% had experienced entanglements (Robbins and Mattila 2001). Decomposed and/or unexamined animals (e.g., carcasses reported but not retrieved or no necropsy performed) represent 'lost data', some of which may relate to human impacts.

Background

As with right whales, human impacts (vessel collisions and entanglements) may be slowing recovery of the humpback whale population. Van der Hoop *et al.* (2013) reviewed 1762 mortalities and serious injuries recorded for 8 species of large whales in the Northwest Atlantic for the 40 years 1970–2009. Of 473 records of humpback whales, cause of death could be attributed for 203. Of the 203, 116 (57%) mortalities were caused by entanglements in fishing gear, and 31 (15%) were attributable to vessel strikes.

Robbins and Mattila (2001) reported that males were more likely to be entangled than females. Annually updated inferences made from scar prevalence and multistate models of GOM humpback whales that (1) younger animals are more likely to become entangled than adults, (2) juvenile scarring rates may be trending up, (3) maybe less than 10% of humpback entanglements are ever reported, and (4) 3 % of the population maybe dying annually as the result of entanglements (Robbins 2009, 2010, 2011, 2012). Humpback whale entanglements also occur in relatively high numbers in Canadian waters. Reports of interactions with fixed fishing gear set for groundfish around Newfoundland averaged 365 annually from 1979 to 1987 (range 174-813). An average of 50 humpback whale entanglements (range 26-66) was reported annually between 1979 and 1988, and 12 of 66 humpback whales entangled in 1988 died (Lien *et al.* 1988). A total of 965 humpbacks was reported entangled in fishing gear in Newfoundland and Labrador from 1979 to 2008 (Benjamins *et al.* 2012). Volgenau *et al.* (1995) reported that in Newfoundland and Labrador, cod traps caused the most entanglements and entanglement mortalities (21%) of humpbacks between 1979 and 1992. They also reported that gillnets were the primary cause of entanglements and entanglement mortalities (20%) of humpbacks in the Gulf of Maine between 1975 and 1990. In more recent times, following the collapse of the cod fishery, groundfish gillnets for other fish species and crab pot lines have been the most common sources of humpback entanglement in Newfoundland. Since the crab pot fishery is primarily an offshore activity on the Grand Banks, these entanglements are hard to respond to and are likely underreported. One humpback whale was reported released alive (status unknown) from a herring weir off Grand Manan in 2009 (H. Koopman, UNC Wilmington, pers. comm.). In US waters, Johnson *et al.* (2005) found 40% of humpback entanglements were in trap/ pot gear and 50% were in gillnet, but sample sizes were small and much uncertainty still exists about the frequency of certain gear types involved in entanglement.

Wiley *et al.* (1995) reported that serious injuries attributable to ship strikes are more common and probably more serious than those from entanglements, but this claim is not supported by more recent analysis. Non-lethal

interactions with gear are extremely common (see Robbins 2010, 2011, 2012) and recent analysis suggests entanglement serious injuries and mortalities are more common than ship strikes (van der Hoop *et al.* 2013). Furthermore, in the NMFS records for 2009 through 2013, there are only 8 reports of serious injuries and mortalities as a result of collision with a vessel and 41 records of injuries (prorated or serious) and mortalities attributed to entanglement. Because it has never been shown that serious injuries and mortalities related to ships or to fisheries interactions are equally detectable, it is unclear as to which human source of mortality is more prevalent. A major aspect of vessel collision that will be cryptic as a serious injury is blunt trauma, where when lethal it is usually undetectable from an external exam (Moore *et al.* 2013). No whale involved in the recorded vessel collisions had been identified as a member of a stock other than the Gulf of Maine stock at the time of this writing (Henry *et al.* 2015).

Fishery-Related Serious Injuries and Mortalities

A description of fisheries is provided in Appendix III. Two mortalities were observed in the pelagic drift gillnet fishery, one in 1993 and the other in 1995. In winter 1993, a juvenile humpback was observed entangled and dead in a pelagic drift gillnet along the 200-m isobath northeast of Cape Hatteras. In early summer 1995, a humpback was entangled and found dead in a pelagic drift gillnet on southwestern Georges Bank. Additional reports of mortality and serious injury, as well as description of total human impacts, are contained in records maintained by NMFS. A number of these records (11 entanglements involving lobster pot/trap gear) from the 1990–1994 period were the basis used to reclassify the lobster fishery (62 FR 33, Jan. 2, 1997). Large whale entanglements are rarely observed during fisheries sampling operations. However, during 2008, 3 humpback whales were observed as incidental bycatch: 2 in gillnet gear (1 no serious injury; 1 undetermined) and 1 in a purse seine (released alive), in 2011 a humpback was caught on an observed gillnet trip (disentangled and released free of gear; Henry *et al.* 2015), and in 2012 there was an observed interaction with a humpback whale in mid-Atlantic gillnet gear (non-serious injury). A recent review (Cassoff *et al.* 2011) describes in detail the types of injuries that baleen whales, including humpbacks, suffer as a result of entanglement in fishing gear.

For this report, the records of dead, injured, and/or entangled humpbacks (found either stranded or at sea) for the period 2009 through 2013 were reviewed. When there was no evidence to the contrary, events were assumed to involve members of the Gulf of Maine stock. While these records are not statistically quantifiable in the same way as observer fishery records, they provide some indication of the minimum frequency of entanglements. Specifically to this stock, if the calculations of Robbins (2011, 2012) are reasonable then the 3% mortality due to entanglement that she calculates equates to a minimum average rate of 25, which is nearly 10 times PBR.

Table 2. Confirmed human-caused mortality and serious injury records of humpback whales (*Megaptera novaeangliae*) where the cause was assigned as either an entanglement (EN) or a vessel strike (VS): 2009–2013^a

Date ^b	Injury Determination	ID	Location ^b	Assigned Cause	Value against PBR ^c	Country ^d	Gear Type ^e	Description
2/8/2009	Mortality	-	Cape Fear, NC	EN	1	XU	NP	Evidence of entanglement at mouthline, peduncle, & pectoral w/ associated hemorrhaging. Emaciated.
2/16/2009	Mortality	-	Nags Head, NC	EN	1	XU	NP	Evidence of entanglement involving anchoring or heavily weighted gear w/ associated hemorrhaging.

2/25/2009	Serious Injury	-	off Sandy Hook, NJ	EN	1	US	NR	Anchored. Disentangled but SI due to deformed body position that did not substantially improve after disentangle ment.
4/9/2009	Prorated Injury	-	off Provincetown MA	EN	0.75	XU	NR	Full configuration unknown.
4/11/2009	Prorated Injury	-	off Gloucester, MA	EN	0.75	XU	NR	Configuration unclear-- unknown if body wrap is loose or constricting.
5/23/2009	Prorated Injury	-	off Provincetown MA	EN	0.75	XU	NR	Full configuration unknown.
6/9/2009	Serious Injury	Inukshuk	off Provincetown MA	EN	1	US	NR	Constricting body wrap.
9/12/2009	Prorated Injury	2008 Calf of Touchdown	off White Island, Nova Scotia	EN	0.75	CN	WE	Swam out of entrapment in weir, but carrying some gear in an unknown configuration.
9/16/2009	Prorated Injury	-	off Halifax, Nova Scotia	EN	0.75	XC	NR	Full configuration unknown.
10/20/2009	Serious Injury	-	off Halifax, Nova Scotia	EN	1	CN	GN	Disentangled, but in poor condition: emaciated, heavy cyamid load, lethargic.
11/20/2009	Prorated Injury	-	off Goat Island, NC	EN	0.75	XU	NR	Full configuration unknown.
3/7/2010	Serious Injury	-	off Ponte Vedra Beach, FL	EN	1	XU	NR	Constricting body & flipper wraps. May have shed some or all of gear, but severe health decline: emaciated,

								heavy cyamid load.
3/13/2010	Mortality	-	Ocean City Inlet, MD	VS	1	US	-	Skull fractures w/ associated hemorrhaging
5/5/2010	Serious Injury	-	off Northampton, VA	EN	1	XU	NR	Wrap around fluke blades near insertion & trailing gear. Gear likely to become constricting as animal grows.
5/8/2010	Mortality	-	off Point Judith, RI	EN	1	US	GN	Evidence of constricting gear w/ associated hemorrhaging. Fluid filled lungs.
5/15/2010	Mortality	-	Hatteras Inlet, NC	EN	1	XU	NP	Live stranding -euthanized. Necrotic infected wounds at base of flukes & chronic abrasions on head.
5/28/2010	Mortality	-	off Martha's Vineyard, MA	EN	1	XU	GU	Evidence of entanglement w/ associated bruising & edema.
6/10/2010	Mortality	-	Jones Beach State Park, NY	VS	1	US	-	Extensive hemorrhage & edema on right dorsal lateral surface.
7/4/2010	Mortality	-	off Ocean City Inlet, MD	VS	1	US	-	Extensive hemorrhage & edema to left lateral area.
7/26/2010	Prorated Injury	-	off Chatham, MA	EN	0.75	XU	NR	Full configuration unknown.
8/13/2010	Serious Injury	-	off Orleans, MA	EN	1	US	PT	Partial disentanglement, but remaining head wrap likely to become

								constricting.
8/20/2010	Serious Injury	Chili	off Provincetown MA	EN	1	XU	NR	Embedded wraps;health decline: thin, moderate cyamids, sloughing skin, fluke discoloration
9/10/2010	Prorated Injury	-	off White Head Island, New Brunswick	EN	0.75	XC	NR	Full configuration unknown.
10/2/2010	Prorated Injury	-	off Provincetown MA	EN	0.75	XU	NR	Full configuration unknown. Unable to confirm if a resight of 8/20/10 event.
11/27/2010	Mortality	-	off Grand Manan Island, New Brunswick	EN	1	XC	NR	Evidence of constricting wraps on fluke, peduncle, & pectoral
12/23/2010	Serious Injury	-	off Port Everglades Inlet, FL	EN	1	XU	NP	Evidence of recent constricting entanglement & severe health decline.
1/7/2011	Serious Injury	-	off Oregon Inlet, NC	EN	1	US	GN	Extensive entanglement w/ netting covering majority of body including head, blowholes, & flukes. Immobile & drifting.
2/1/2011	Serious Injury	EKG	off Bar Harbor, ME	EN	1	US	NR	Anchored. Cuts were made to gear but whale remained anchored.
3/7/2011	Mortality	-	Thorofare Bay, NC	VS	1	US	-	Live stranded w/ 8 deep lacerations across back. Euthanized.

4/11/2011	Prorated Injury	-	off Rockport, MA	EN	0.75	XU	NR	Full configuration unknown.
5/5/2011	Mortality	-	Little Compton, RI	VS	1	US	-	Hemorrhaging at left jaw associated w/ blunt trauma.
5/27/2011	Mortality	-	Island Beach State Park, NJ	VS	1	US	-	5 broken vertebral processes along left side w/ associated hemorrhaging.
5/30/2011	Prorated Injury	-	off Orleans, MA	EN	0.75	XU	NR	Full configuration unknown.
7/2/2011	Serious Injury	-	off Provincetown MA	EN	1	XU	NP	Young whale. Missing flukes attributed to chronic entanglement. Laceration due to VS appears minor. Significant health decline: emaciated, swimming by use of pectorals only
7/9/2011	Prorated Injury	-	off Monomoy Island, MA	EN	0.75	XU	NR	Full configuration unknown.
7/10/2011	Prorated Injury	-	off Monomoy Island, MA	EN	0.75	XU	NR	Report of two entangled whales but could not confirm that both were entangled. Full configuration unknown.
7/21/2011	Prorated Injury	-	off Oregon Inlet, NC	EN	0.75	XU	NR	Full configuration unknown.
10/10/2011	Serious Injury	Clutter	off Grand Manan Island, New Brunswick	EN	1	XC	NR	Embedded wraps at fluke insertion.
4/29/2012	Serious Injury	-	off Chatham, MA	EN	1	US	NR	SI based on description of body position which indicates

								anchored
7/29/2012	Serious Injury	-	off Gloucester, MA	EN	1	XU	NR	Calf w/ line cutting into peduncle
8/4/2012	Serious Injury	Aphid	off Provincetown MA	EN	1	XU	NR	Line exiting both sides of mouth, under flippers, twisting together aft of the dorsal fin & trailing 75 ft past flukes; no wraps. Health decline: thin w/ graying skin.
8/21/2012	Prorated Injury	2011 Calf of Wizard	off Provincetown MA	EN	0.75	XU	MF	Full configuration unknown
8/24/2012	Serious Injury	Forceps	off Provincetown MA	EN	1	US	NR	Closed, possibly weighted, bridle w/ large tangle of line just above left eye. SI due to odd behavior & apparent difficulty staying at the surface.
04/03/2013	Mortality	-	off Ft Story, VA	VS	1	US	-	Fractured orbitals & ribs w/ associated bruising
09/13/2013	Mortality	-	York River, VA	VS	1	US	-	6 lacerations penetrate into muscle w/ associated hemorrhaging
09/16/2013	Prorated Injury	-	off Chatham, MA	EN	0.75	XU	NR	Partial disentanglement; original & final configurations unknown

09/28/2013	Mortality	-	off Saltaire, NY	EN	1	XU	GU	Embedded line in mouth w/ associated hemorrhaging & necrosis; evidence of constriction at pectorals, peduncle & fluke w/ associated hemorrhaging; emaciated
10/01/2013	Mortality	-	Buzzards Bay, MA	EN	1	US	NP	Evidence of underwater entrapment & subsequent drowning.
10/04/2013	Serious Injury	-	off Chatham, MA	EN	1	XU	NR	Full configuration unknown, but evidence of health decline: emaciation & pale skin
Five-year averages		Shipstrike (US/CN/XU/XC)			1.60 (1.60/ 0.00/ 0.00/ 0.00)			
		Entanglement (US/CN/XU/XC)			7.4 (1.8/ 0.35/ 4.55/ 0.70)			
a. For more details on events please see Henry <i>et al.</i> 2015.								
b. The date sighted and location provided in the table are not necessarily when or where the serious injury or mortality occurred; rather, this information indicates when and where the whale was first reported beached, entangled, or injured.								
c. Mortality events are counted as 1 against PBR. Serious injury events have been evaluated using NMFS guidelines (NOAA 2012)								
d. CN=Canada, US=United States, XC=Unassigned 1st sight in CN, XU=Unassigned 1st sight in US								
e. H=hook, GN=gillnet, GU=gear unidentifiable, MF=monofilament, NP=none present, NR=none recovered/received, PT=pot/trap, WE=weir								

Other Mortality

Between November 1987 and January 1988, at least 14 humpback whales died after consuming Atlantic mackerel containing a dinoflagellate saxitoxin (Geraci *et al.* 1989). The whales subsequently stranded or were recovered in the vicinity of Cape Cod Bay and Nantucket Sound, and it is highly likely that other unrecorded mortalities occurred during this event. During the first six months of 1990, seven dead juvenile (7.6 to 9.1 m long) humpback whales stranded between North Carolina and New Jersey. The significance of these strandings is unknown.

Between July and September 2003, an Unusual Mortality Event (UME) that included 16 humpback whales was invoked in offshore waters of coastal New England and the Gulf of Maine. Biotxin analyses of samples taken from some of these whales found saxitoxin at very low/questionable levels and domoic acid at low levels, but neither were adequately documented and therefore no definitive conclusions could be drawn. Seven humpback whales were considered part of a large whale UME in New England in 2005. Twenty-one dead humpback whales found between 10 July and 31 December 2006 triggered a humpback whale UME declaration. Causes of these UME events have not been determined.

STATUS OF STOCK

NMFS has conducted a global humpback whale status review (Bettridge *et al.* 2015), and recently proposed

revising the ESA listing of humpback whales (80 FR 22303, April 21, 2015). Although recent estimates of abundance indicate a stable or growing humpback whale population, the stock may be below OSP in the U.S. Atlantic EEZ. The detected level of U.S. fishery-caused mortality and serious injury derived from the available records, which is likely biased low, is more than 10% of the calculated PBR and, therefore, cannot be considered to be insignificant or approaching a zero mortality and serious injury rate. This is a strategic stock because the average annual human-related mortality and serious injury exceeds PBR, and because the North Atlantic humpback whale is an endangered species.

REFERENCES CITED

- Balcomb, K.C. and G. Nichols. 1982. Humpback whale censuses in the West Indies. Rep. Int. Whal. Comm. 32: 401-406.
- Barco, S.G., W.A. McLellan, J.M. Allen, R.A. Asmutis-Silvia, R. Mallon-Day, E.M. Meagher, D.A. Pabst, J. Robbins, R.E. Seton, W.M. Swingle, M.T. Weinrich, and P.J. Clapham. 2002. Population identity of humpback whales (*Megaptera novaeangliae*) in the waters of the U.S. mid-Atlantic states. J. Cetacean Res. Manage. 4(2): 135-141.
- Barlow, J. and P.J. Clapham. 1997. A new birth-interval approach to estimating demographic parameters of humpback whales. Ecology 78: 535-546.
- Barlow, J., S.L. Swartz, T.C. Eagle, and P.R. Wade. 1995. U.S. marine mammal stock assessments: Guidelines for preparation, background, and a summary of the 1995 assessments. NOAA Tech. Memo. NMFS-OPR-6. 73 pp.
- Benjamins, S., W. Ledwell, J. Huntington, and A.R. Davidson. 2012. Assessing changes in numbers and distribution of large whale entanglements in Newfoundland and Labrador, Canada. Mar. Mamm. Sci. 28(3):579-601.
- Bettridge, S., C.S. Baker, J. Barlow, P.J. Clapham, M. Ford, D. Gouveia, D.K. Matila, R.M. Pace III., P.E. Rosel, G.K. Silber, and P.R. Wade. 2015. Status Review of the Humpback Whale (*Megaptera novaeangliae*) under the Endangered Species Act. NOAA Technical Memorandum NMFS. NOAA-TM-NMFS-SWFSC-540 NMFS, Southwest Fisheries Science Center, California.
- Brandão, A., D.S. Butterworth, and M.R. Brown. 2000. Maximum possible humpback whale increase rates as a function of biological parameter values. J. Cetacean Res. Manage. 2 (supplement): 192-193.
- Cassoff, R., K. Moore, W. McLellan, S. Barco, D. Rotstein, and M. Moore. 2011. Lethal entanglement in baleen whales. Dis. Aquat. Org. 96:175-185.
- Christensen, I., T. Haug, and N. Øien. 1992. Seasonal distribution, exploitation and present abundance of stocks of large baleen whales (Mysticeti) and sperm whales (*Physeter macrocephalus*) in Norwegian and adjacent waters. ICES J. Mar. Sci. 49: 341-355.
- Clapham, P.J., L.S. Baraff, C.A. Carlson, M.A. Christian, D.K. Mattila, C.A. Mayo, M.A. Murphy, and S. Pittman. 1993. Seasonal occurrence and annual return of humpback whales, *Megaptera novaeangliae*, in the southern Gulf of Maine. Can. J. Zool. 71: 440-443.
- Clapham, P.J., J. Barlow, M. Bessinger, T. Cole, D. Mattila, R. Pace, D. Palka, J. Robbins, and R. Seton 2003. Abundance and demographic parameters of humpback whales from the Gulf of Maine, and stock definition relative to the Scotian Shelf. J. Cetacean Res. Manage. 5(1): 13-22.
- Clapham, P.J., M.C. Bérubé, and D.K. Mattila. 1995. Sex ratio of the Gulf of Maine humpback whale population. Mar. Mamm. Sci. 11: 227-231.
- Clapham, P.J. and C.A. Mayo. 1987. Reproduction and recruitment of individually identified humpback whales, *Megaptera novaeangliae*, observed in Massachusetts Bay, 1979-1985. Can. J. Zool. 65: 2853-2863.
- Clapham, P.J., J. Robbins, M. Brown, P. Wade and K. Findlay. 2001. A note on plausible rates of population growth for humpback whales. J. Cetacean Res. Manage. 3 (suppl.): 196-197.
- Fogarty, M.J., E.B. Cohen, W.L. Michaels, and W.W. Morse. 1991. Predation and the regulation of sand lance populations: An exploratory analysis. ICES Mar. Sci. Symp. 193: 120-124.
- Friedlaender, A., E. Hazen, D. Nowacek, P. Halpin, C. Ware, M. Weinrich, T. Hurst, and D. Wiley. 2009. Diel changes in humpback whale *Megaptera novaeangliae* feeding behavior in response to sand lance *Ammodytes* spp. behavior and distribution. Mar. Ecol. Prog. Ser. 395:91-100.
- Geraci, J.R., D.M. Anderson, R.J. Timperi, D.J.S. Aubin, G.A. Early, J.H. Prescott, and C.A. Mayo. 1989. Humpback whales (*Megaptera novaeangliae*) fatally poisoned by dinoflagellate toxins. Can. J. Fish. Aquat. Sci. 46: 1895-1898.
- [Henry, A.G., T.V.N. Cole, L. Hall, W. Ledwell, D. Morin and A. Reid. 2015. Mortality and Serious Injury determinations for baleen whale stocks along the Gulf of Mexico, United States East Coast and Atlantic Canadian Provinces, 2009–2013, NEFSC Reference Document 15-10. 45 pp.](#)

- IWC 2002. Report of the Scientific Committee. Annex H: Report of the Sub-committee on the Comprehensive Assessment of North Atlantic humpback whales. *J. Cetacean Res. Manage.* 4 (supplement): 230-260.
- Johnson, A., G. Salvador, J. Kenney, J. Robbins, S. Kraus, S. Landry, and P. Clapham .2005. Fishing gear involved in entanglements of right and humpback whales. *Mar. Mamm.Sci.*21(4):635–645.
- Katona, S.K. and J.A. Beard. 1990. Population size, migrations, and feeding aggregations of the humpback whale (*Megaptera novaeangliae*) in the western North Atlantic Ocean. *Rep. Int. Whal. Comm. (Special Issue)* 12: 295-306.
- Laake, J.L. and D.L. Borchers. 2004. Methods for incomplete detection at distance zero. Pages 108-189 in: *Advanced distance sampling*, edited by S.T. Buckland, D.R. Andersen, K.P. Burnham, J.L. Laake, and L. Thomas. Oxford University Press, New York.
- Larsen, A.H., J. Sigurjónsson, N. Øien, G. Vikingsson, and P.J. Palsbøll. 1996. Population genetic analysis of mitochondrial and nuclear genetic loci in skin biopsies collected from central and northeastern North Atlantic humpback whales (*Megaptera novaeangliae*): population identity and migratory destinations. *Proc. R. Soc. London, B* 263: 1611-1618.
- Levenson, C. and W.T. Leapley. 1978. Distribution of humpback whales (*Megaptera novaeangliae*) in the Caribbean determined by a rapid acoustic method. *J. Fish. Res. Board Can.* 35: 1150-1152.
- Lien, J., W. Ledwell, and J. Naven. 1988. Incidental entrapment in inshore fishing gear during 1988: A preliminary report to the Newfoundland and Labrador Department of Fisheries and Oceans. St. John's, Newfoundland, Canada 15 pp.
- Mattila, D.K. and P.J. Clapham. 1989. Humpback whales and other cetaceans on Virgin Bank and in the northern Leeward Islands, 1985 and 1986. *Can. J. Zool.* 67: 2201-2211.
- Mattila, D.K., P.J. Clapham, S.K. Katona, and G.S. Stone. 1989. Population composition of humpback whales on Silver Bank. *Can. J. Zool.* 67: 281-285.
- Moore, M., J. van der Hoop, S. Barco, A. Costidis, F. Gulland, P. Jepson, K. Moore, S. Raverty, and W. McLellan. 2013. Criteria and case definitions for serious injury and death of pinnipeds and cetaceans caused by anthropogenic trauma. *Dis.of Aq. Org.* 103(3):229–264.
- Palka, D.L. 2012. Cetacean abundance estimates in US northwestern Atlantic Ocean waters from summer 2011 line transect survey. *Northeast Fish. Sci. Cent. Ref. Doc.* 12-29. 37 pp.
- Palka, D.L. 2006. Summer abundance estimates of cetaceans in US North Atlantic Navy Operating Areas. *Northeast Fish. Sci. Cent. Ref. Doc.* 06-03. 41 pp.
<http://www.nefsc.noaa.gov/nefsc/publications/crd/crd0603/crd0603.pdf>
- Palsbøll, P.J., J. Allen, T.H. Anderson, M. Bérubé, P.J. Clapham, T.P. Feddersen, N. Friday, P. Hammond, H. Jørgensen, S.K. Katona, A.H. Larsen, F. Larsen, J. Lien, D.K. Mattila, F.B. Nygaard, J. Robbins, R. Spomer, R. Sears, J. Sigurjónsson, T.D. Smith, P.T. Stevick, G. Vikingsson, and N. Øien. 2001. Stock structure and composition of the North Atlantic humpback whale, *Megaptera novaeangliae*. International Whaling Commission Scientific Committee, Cambridge, UK. SC/53/NAH11.
- Palsbøll, P.J., J. Allen, M. Berube, P. Clapham, T. Feddersen, P. Hammond, R. Hudson, H. Jorgensen, S. Katona, A.H. Larsen, F. Larsen, J. Lien, D. Mattila, J. Sigurjónsson, R. Sears, T. Smith, R. Spomer, P. Stevick, and N. Oien. 1997. Genetic tagging of humpback whales. *Nature* 388: 767-769.
- Palsbøll, P.J., P.J. Clapham, D.K. Mattila, F. Larsen, R. Sears, H.R. Siegismund, J. Sigurjónsson, O. Vásquez, and P. Arctander. 1995. Distribution of mtDNA haplotypes in North Atlantic humpback whales: the influence of behavior on population structure. *Mar. Ecol. Prog. Ser.* 116: 1-10.
- Paquet, D., C. Haycock, and H. Whitehead. 1997. Numbers and seasonal occurrence of humpback whales (*Megaptera novaeangliae*) off Brier Island, Nova Scotia. *Can. Field-Nat.* 111: 548-552.
- Payne, P.M., J.R. Nicholas, L. O'Brien, and K.D. Powers. 1986. The distribution of the humpback whale, *Megaptera novaeangliae*, on Georges Bank and in the Gulf of Maine in relation to densities of the sand eel, *Ammodytes americanus*. *Fish. Bull.* 84: 271-277.
- Payne, P.M., D.N. Wiley, S.B. Young, S. Pittman, P.J. Clapham, and J.W. Jossi. 1990. Recent fluctuations in the abundance of baleen whales in the southern Gulf of Maine in relation to changes in selected prey. *Fish. Bull.* 88: 687-696.
- Price, W.S. 1985. Whaling in the Caribbean: historical perspective and update. *Rep. Int. Whal. Comm.* 35: 413-420.
- Reiner, F., M.E.D. Santos, and F.W. Wenzel. 1996. Cetaceans of the Cape Verde archipelago. *Mar. Mamm. Sci.* 12: 434-443.
- Robbins, J. 2007. Structure and dynamics of the Gulf of Maine humpback whale population. Ph.D. thesis. University of St. Andrews, Aberdeen, Scotland.

- Robbins, J. 2009. Scar based inference into Gulf of Maine Humpback whale Entanglement: 2003-2006. Report to the Northeast Fisheries Science Center, National Marine Fisheries Service, Woods Hole, Massachusetts, USA. <http://www.nefsc.noaa.gov/psb/docs/HUWHScarring%28Robbins2009%29.pdf>
- Robbins, J. 2010. Scar-based inference into Gulf of Maine Humpback whale Entanglement: 2008. Report to the Northeast Fisheries Science Center, National Marine Fisheries Service, Woods Hole, Massachusetts, USA.
- Robbins, J. 2011. Scar-based inference into Gulf of Maine Humpback whale Entanglement: 2009. Report to the Northeast Fisheries Science Center, National Marine Fisheries Service, Woods Hole, Massachusetts, USA. <http://www.nefsc.noaa.gov/psb/docs/HUWHScarring%28Robbins2011%29.pdf>
- Robbins, J. 2012. Scar-based inference into Gulf of Maine Humpback whale Entanglement: 2010. Report to the Northeast Fisheries Science Center, National Marine Fisheries Service, Woods Hole, Massachusetts, USA. <http://www.nefsc.noaa.gov/psb/docs/HUWHScarring%28Robbins2012%29.pdf>
- Robbins, J. and D.K. Mattila. 2001. Monitoring entanglements of humpback whales (*Megaptera novaeangliae*) in the Gulf of Maine on the basis of caudal peduncle scarring. International Whaling Commission Scientific Committee, Cambridge, UK. SC/53/NAH25.
- Smith, T.D., J. Allen, P.J. Clapham, P.S. Hammond, S. Katona, F. Larsen, J. Lien, D. Mattila, P.J. Palsboll, J. Sigurjónsson, P.T. Stevick, and N. Øien. 1999. An ocean-basin-wide mark-recapture study of the North Atlantic humpback whale (*Megaptera novaeangliae*). Mar. Mamm. Sci. 15(1): 1-32.
- Smith, T.D. and D.G. Pike. 2009. The enigmatic humpback whale. NAMMCO Sci. Publ. 7:161-178.
- Stevick, P., N. Øien, and D.K. Mattila. 1998. Migration of a humpback whale between Norway and the West Indies. Mar. Mamm. Sci. 14: 162-166.
- Stevick, P.T., J. Allen, P.J. Clapham, N. Friday, S.K. Katona, F. Larsen, J. Lien, D.K. Mattila, P.J. Palsbøll, J. Sigurjónsson, T.D. Smith, N. Øien, and P.S. Hammond. 2003. North Atlantic humpback whale abundance and rate of increase four decades after protection from whaling. Mar. Ecol. Prog. Ser. 258: 263-273.
- Stevick, P.T., J. Allen, P.J. Clapham, S.K. Katona, F. Larsen, J. Lien, D.K. Mattila, P.J. Palsbøll, R. Sears, J. Sigurjónsson, T.D. Smith, G. Vikingsson, N. Øien and P.S. Hammond. 2006. Population spatial structuring on the feeding grounds in North Atlantic humpback whales (*Megaptera novaeangliae*). J. Zool. 270: 244-255.
- Swingle, W.M., S.G. Barco, T.D. Pitchford, W.A. McLellan, and D.A. Pabst 1993. Appearance of juvenile humpback whales feeding in the nearshore waters of Virginia. Mar. Mamm. Sci. 9: 309-315.
- Thomas L, J.L. Laake, E. Rexstad, S. Strindberg, F.F.C. Marques, S.T. Buckland, D.L. Borchers, D.R. Anderson, K.P. Burnham, M.L. Burt, S.L. Hedley, J.H. Pollard, J.R.B. Bishop, and T.A. Marques. 2009. Distance 6.0. Release 2. [Internet]. University of St. Andrews (UK): Research Unit for Wildlife Population Assessment. Available from: <http://www.ruwpa.st-and.ac.uk/distance/>
- van Der Hoop, J.M., M.J. Moore, S.G. Barco, T.V.N. Cole, P.-Y. Daoust, A.G. Henry, D.F. McAlpine, W.A. McLellan, T. Wimmer, and A.R. Solow. 2013. Assessment of management to mitigate anthropogenic effects on large whales. Cons. Biol. 27(1):121-133.
- Volgenau, L., S.D. Kraus, and J. Lien. 1995. The impact of entanglements on two substocks of the western North Atlantic humpback whale, *Megaptera novaeangliae*. Can. J. Zool. 73: 1689-1698.
- Vu, E., D. Risch, C. Clark, S. Gaylord, L. Hatch, M. Thompson, D. Wiley, and S. Van Parijs. 2012. Humpback whale song occurs extensively on feeding grounds in the western North Atlantic Ocean. Aq. Biol. 14(2): 175-183.
- Wade, P.R. and R.P. Angliss 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Waring, G.T., D.L. Palka, P.J. Clapham, S. Swartz, M.C. Rossman, T.V.N. Cole, K.D. Bisack, and L.J. Hansen, eds. 1999. U.S. Atlantic marine mammal stock assessments - 1998. NOAA Tech. Memo. NMFS-NE-116. 182 pp.
- Waring, G.T., J.M. Quintal, and S.L. Swartz, eds. 2000. U.S. Atlantic and Gulf of Mexico marine mammal stock assessments - 2000. NOAA Tech. Memo. NMFS-NE-162. ix +197 pp. +3 app.
- Weinrich, M., M. Martin, R. Griffiths, J. Bove, and M. Schilling. 1997. A shift in distribution of humpback whales, *Megaptera novaeangliae*, in response to prey in southern Gulf of Maine. Fish. Bull. 95:826-836.
- Wenzel, F.W., J. Allen, S. Berrow, C.J. Hazevoet, B. Jann, R.E. Seton, L. Steiner, P. Stevick, P.L. Suárez, and P. Whooley. 2009. Current knowledge on the distribution and relative abundance of humpback whales (*Megaptera novaeangliae*) off the Cape Verde Islands, eastern North Atlantic. Aquat. Mamm. 35: 502-510.
- Wiley, D.N., R.A. Asmutis, T.D. Pitchford, and D.P. Gannon 1995. Stranding and mortality of humpback whales, *Megaptera novaeangliae*, in the mid-Atlantic and southeast United States, 1985-1992. Fish. Bull. 93: 196-205.

- Winn, H.E., R.K. Edel, and A.G. Taruski. 1975. Population estimate of the humpback whale (*Megaptera novaeangliae*) in the West Indies by visual and acoustic techniques. J. Fish. Res. Board Can. 32: 499-506.
- Zerbini, A.N., P.J. Clapham, and P.R. Wade. 2010. Assessing plausible rates of population growth in humpback whales from life-history data. Mar. Biol. 157(6):1225–1236.

