

NARWHAL (*Monodon monoceros*): Unidentified Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Narwhals typically inhabit waters of the Arctic Ocean. They are common in the waters of Nunavut, Canada, west Greenland, and in the European Arctic; however, they rarely occur in the East Siberian, Bering, Chukchi, and Beaufort Seas (COSEWIC 2004). The three recognized populations of narwhals are based on summer distribution: Baffin Bay, Hudson Bay, and east Greenland (DFO 1998a, 1998b; COSEWIC 2004). The Baffin Bay population of narwhals summers in the waters of West Greenland and the Canadian High Arctic and overwinters in Baffin Bay and Davis Strait (Koski and Davis 1994; Dietz et al. 2001; Heide-Jørgensen et al. 2003). Narwhals from the northwest Hudson Bay population are thought to overwinter in eastern Hudson Strait (Richard 1991). The east Greenland population is believed to winter in the pack ice between eastern Greenland and Svalbard (Dietz et al. 1994). The amount of interchange between these populations is unknown; populations are defined for management purposes, and these designated populations may actually consist of several populations (COSEWIC 2004). Population definition based on molecular genetics studies of narwhals remains unresolved at this time due to extremely low genetic variability within and among management stocks (Palsbøll et al. 1997; de March et al. 2001, 2003).

Local observations and traditional ecological knowledge are the primary source for observation data of narwhals in Alaska waters, dating back to the 1800s (Bee and Hall 1956, Geist et al. 1960, Noongwook et al. 2007, George and Suydam unpubl. ms.). The earliest record dates back to 1874, with most of the occasional sightings occurring around the area east of Point Barrow (Scammon 1874, Ray and Murdoch 1885, Turner 1886, Nelson and True 1887, Murdoch 1898, MacFarlane 1905, Dufresne 1946, Anderson 1947, Bee and Hall 1956, Geist et al. 1960). Narwhal occurrences are reported in Bee and Hall (1956) from Pt. Barrow to the Colville River Delta. Ljungblad et al. (1983) reported on a sighting of two male narwhals that occurred northwest of King Island in the Bering Sea, just south of the Bering Strait, during a systematic scientific survey. Sightings have occurred in Russian waters of the northern Chukchi Sea in Russian waters (Reeves and Tracey 1980, Yablokov and Bel'kovich 1968). George and Suydam (unpubl. ms.) summarized observations from Alaska Native hunters during eight sighting events of narwhals in the Chukchi and Beaufort Seas between 1989 and 2008. Of these records, seven were sightings of live animals totaling 11-12 individuals; one record was a report of a beach cast narwhal tusk at Cape Sabine. Four of the seven sightings of live animals consisted of mixed groups of beluga and narwhals (George and Suydam unpubl. ms.). It is believed that these incidental sightings of narwhals occurring in the Beaufort, Chukchi, and Bering seas are whales from the Baffin Bay population that are known to move into the Canadian Arctic Archipelago and as far north and west as ice conditions will permit (COSEWIC 2004).

Several specimens of narwhals collected in Alaska have been documented. Huey (1952) reported on a specimen collected near Cape Halkett, Harrison Bay, at the mouth of the Colville River. Three additional specimen records from various locations were documented in Geist et al. (1960); one specimen was found dead on the beach of Kiwalik Bay (Kotzebue Sound), another was initially sighted alive at the mouth of the Caribou River in Nelson Lagoon on the Alaska Peninsula but later died, and a third specimen of a narwhal tusk was found on the beach at Wainwright. Murie (1936) reported on a single tusk that was found on a sandbar at Cape Chibukak, St. Lawrence Island.



Figure 20. Potential distribution of narwhals in Arctic waters based on extralimital sightings and strandings (George and Suydam, unpubl. ms., Reeves and Tracey 1980, COSEWIC 2004).

Narwhal in Alaska are thought to originate from a Canadian population, but there is no available method to verify this. There are insufficient data to apply the phylogeographic approach to stock structure (Dizon et al. 1992) for narwhal.

POPULATION SIZE

Reliable estimates of abundance for narwhal in Alaska are currently unavailable.

Minimum Population Estimate

At this time, it is not possible to produce a reliable minimum population estimate (N_{MIN}) for this stock, as current estimates of abundance are unavailable.

Current Population Trend

At present, reliable data on trends in population abundance are unavailable.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

A reliable estimate of the maximum net productivity rate is currently unavailable for narwhals in Alaska. Hence, until additional data become available, it is recommended that the cetacean maximum theoretical net productivity rate (R_{MAX}) of 4% be employed (Wade and Angliss 1997).

POTENTIAL BIOLOGICAL REMOVAL

Under the 1994 reauthorized Marine Mammal Protection Act (MMPA), the potential biological removal (PBR) is defined as the product of the minimum population estimate, one-half the maximum theoretical net productivity rate, and a recovery factor: $PBR = N_{\text{MIN}} \times 0.5R_{\text{MAX}} \times F_R$. The recovery factor (F_R) for these stocks is 0.5, the value for cetacean stocks with unknown population status (Wade and Angliss 1997). However, in the absence of a reliable estimate of minimum abundance, the PBR for this stock is unknown.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

New Serious Injury Guidelines

NMFS updated its serious injury designation and reporting process, which uses guidance from previous serious injury workshops, expert opinion, and analysis of historic injury cases to develop new criteria for distinguishing serious from non-serious injury (Angliss and DeMaster 1998, Andersen *et al.* 2008, NOAA 2012). NMFS defines serious injury as an “*injury that is more likely than not to result in mortality*”. Injury determinations for stock assessments revised in 2013 or later incorporate the new serious injury guidelines, based on the most recent 5-year period for which data are available.

Fisheries Information

There are no U. S. commercial fisheries operating within the range of the narwhals in Alaska. There are no observer program records of narwhal mortalities incidental to commercial fisheries in Alaska. The estimated annual mortality rate incidental to commercial fisheries is zero.

Subsistence/Native Harvest Information

There is no known subsistence harvest of narwhals by Alaska Natives.

STATUS OF STOCK

Narwhals are not designated as “depleted” under the MMPA or listed as “threatened” or “endangered” under the Endangered Species Act. Reliable estimates of the minimum population, population trend, PBR, and status of the stock relative to its Optimum Sustainable Population size are currently not available. There are no federal or state commercial fisheries operating in the marine waters of the Arctic, and there are no reports of serious injury or mortality of narwhals in Alaska, so the level of serious injury and mortality is considered insignificant and approaching zero. The estimated annual rate of human-caused mortality and serious injury is believed to be zero for this stock. Thus, the unidentified stock of narwhals is not classified as strategic.

CITATIONS

- Andersen, M. S., K. A. Forney, T. V. N. Cole, T. Eagle, R. Angliss, K. Long, L. Barre, L. Van Atta, D. Borggaard, T. Rowles, B. Norberg, J. Whaley, and L. Engleby. 2008. Differentiating Serious and Non-Serious Injury of Marine Mammals: Report of the Serious Injury Technical Workshop, 10-13 September 2007, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-39. 94 p.
- Anderson, R. M. 1947. Catalogue of Canadian recent mammals. Bull. Natl. Mus. Canada, Biol. Ser., No. 31: vi + 238.
- Angliss, R. P. and D. P. DeMaster. 1998. Differentiating Serious and Non-Serious Injury of Marine Mammals Taken Incidental to Commercial Fishing Operations. NOAA Tech Memo. NMFS-OPR-13, 48 p.
- Bee, J. W. and E. R. Hall. 1956. Mammals of northern Alaska on the Arctic slope. Univ. Kansas Mus. Nat. Hist., Misc. Publ. No. 8, 309 pp.
- COSEWIC 2004. COSEWIC assessment and update status report on the narwhal, *Monodon monoceros*, in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. vii + 50 pp. (www.sararegistry.gc.ca/status/status_e.cfm).
- de March, B.G.E., L.D. Maiers, and D. Tenkula. 2001. A preliminary analysis of the molecular genetics of narwhal (*Monodon monoceros*) samples collected from Canadian and adjacent waters from 1982 to 2000. Canada/Greenland Joint Commission on the Management and Conservation of Narwhal and Beluga (JCNB), Scientific Working Group, Quqetarsuaq, Greenland, May 9-13, 2001. Document No. SWG-2001-10.
- de March, B.G.E., D.A. Tenkula, and L.D. Postma. 2003. Molecular genetics of narwhal (*Monodon monoceros*) from Canada and West Greenland (1982-2001). Canada Department of Fisheries and Oceans, Canadian Science Advisory Secretariat Research Document 2003/080: 23 p.
- DFO [Canada. Department of Fisheries and Oceans]. 1998a. Hudson Bay narwhal. Canada Department of Fisheries and Oceans, Central and Arctic Region, DFO Sci. Stock Status Rep. E5-44: 5 p.
- DFO [Canada. Department of Fisheries and Oceans]. 1998b. Baffin Bay narwhal. Canada Department of Fisheries and Oceans, Central and Arctic Region, DFO Sci. Stock Status Rep. E5-43: 5 p.
- Dietz, R., M.P. Heide-Jørgensen, P.R. Richard, and M. Acquarone. 2001. Summer and fall movements of narwhals (*Monodon monoceros*) from northeastern Baffin Island towards northern Davis Strait. Arctic 54: 244-261.
- Dietz, R., M.P. Heide-Jørgensen, E. Born, and C.M. Glahder. 1994. Occurrence of narwhals (*Monodon monoceros*) and white whales (*Delphinapterus leucas*) in East Greenland. Medd. Grøn. Biosci. 39: 69-86.
- Dizon, A. E., C. Lockyer, W. F. Perrin, D. P. DeMaster, and J. Sisson. 1992. Rethinking the stock concept: a phylogeographic approach. Conserv. Biol. 6:24-36.
- Dufresne, F. 1946. Alaska's animals and fishes. New York: A. S. Barnes. xviii + 297 pp.
- Geist, O. W., J. L. Buckley, R. H. Manville. 1960. Alaskan records of the narwhal. J. Mammal., Vol. 41, No. 2: 250-253.
- George, J. C. and Suydam, R. unpubl. ms. Recent observations of narwhal in the Chukchi and Beaufort Seas by local hunters. 13 January 2009. 3 pp. (Available from D. Allen, National Marine Mammal Laboratory, Alaska Fisheries Science Center, 7600 Sand Point Way, NE, Seattle, WA 98115).
- Heide-Jørgensen, M.P., R. Dietz, K.L. Laidre, P.R. Richard, J. Orr, and H.C. Schmidt. 2003. The migratory behaviour of narwhals (*Monodon monoceros*). Can. J. Zool. 81: 1298-1305.
- Huey, L. M. 1952. An Alaskan record of the narwhal. J. Mammal. 33:496.
- Koski, W.R. and R.A. Davis. 1994. Distribution and numbers of narwhals (*Monodon monoceros*) in Baffin Bay and Davis Strait. Medd. Grøn. Biosci. 39: 15-40.
- Ljungblad, D. K., S. E. Moore, and D. R. Van Schoik. 1983. Aerial surveys of endangered whales in the Beaufort, Eastern Chukchi and Northern Bering Seas, 1982 NOSC Technical Document 605. 110 pp plus appx.
- MacFarlane, R. 1905. Notes on mammals collected and observed in the northern Mackenzie River District, Northwest Territories of Canada. Proc. U. S. Nat. Mus., 28: 673-764.
- Murdoch, J. 1898. The animals known to the Eskimos of Northwestern Alaska. Amer. Nat., 32: 719-734.
- Murie, O. J. 1936. Notes on the mammals of St. Lawrence Island, Alaska. In Archaeological Excavations at Kukulik, St. Lawrence Island, Alaska. Univ. Alaska, Misc. Publ., 2: 337-346.
- Nelson, E. W. and F. W. True. 1887. Mammals of northern Alaska. Pt. 2 in Report upon natural history collections made in Alaska between the years 1877 and 1881 by Edward W. Nelson. Arctic Publ. No. 3, Signal Service, U. S. Army; pp. 227-293.

- Noongwook, G., The Native Village of Savoonga, The Native Village of Gambell, Huntington, H.P., and George, J.C. 2007. Traditional knowledge of the bowhead whale (*Balaena mysticetus*) around St. Lawrence Island, Alaska. *Arctic* 60 (1): 47-54.
- NOAA. 2012. Federal Register 77:3233. National Policy for Distinguishing Serious From Non-Serious Injuries of Marine Mammals. <http://www.nmfs.noaa.gov/op/pds/documents/02/238/02-238-01.pdf>.
- Palsbøll, P.J., M.P. Heide-Jørgensen, and R. Dietz. 1997. Genetic studies of narwhals, *Monodon monoceros*, from West and East Greenland. *Heredity* 78: 284-292.
- Ray, P. H. and J. Murdoch. 1885. Report of the International Polar Expedition to Point Barrow, Alaska. Washington, 695 pp.
- Reeves, R. R. and Tracey, S. 1980. *Monodon monoceros*. *Mamm. Species*, 127:1-7, 5 figs.
- Richard, P. 1991. Abundance and distribution of narwhals (*Monodon monoceros*) in northern Hudson Bay. *Can. J. Fish. Aquat. Sci.* 48: 276-283.
- Scammon, C. M. 1874. The marine mammals of the northwestern coast of North America, described and illustrated: together with an account of the American whale fishery. New York: G. P. Putnam's Sons. 319 pp.
- Turner, L. M. 1886. Contributions to the natural history of Alaska; results of investigations made chiefly in the Yukon District and the Aleutian Islands. *Arctic Publ. No. 2*, Signal Service, U. S. Army; pp. 1-226.
- Wade, P. R., and R. Angliss. 1997. Guidelines for assessing marine mammal stocks: report of the GAMMS workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Yablokov A. V. and V. M. Bel'kovich. 1968. Cetaceans of the arctic; their proper utilization and conservation. *Probl. of the North, Nat. Res. Council, Ottawa*, 11:199-218.