

## MESOPLDONT BEAKED WHALES (*Mesoplodon* spp.): California/Oregon/Washington Stocks


### STOCK DEFINITION AND GEOGRAPHIC RANGE

Mesoplodont beaked whales are distributed throughout deep waters and along the continental slopes of the North Pacific Ocean. The six species known to occur in this region are: Blainville's beaked whale (*M. densirostris*), Perrin's beaked whale (*M. perrini*), Lesser beaked whale (*M. peruvianus*), Stejneger's beaked whale (*M. stejnegeri*), Ginkgo-toothed beaked whale (*M. ginkgodens*), and Hubbs' beaked whale (*M. carlhubbsi*) (Mead 1989, Henshaw *et al.* 1997, Dalebout *et al.* 2002, MacLeod *et al.* 2006). Based on bycatch and stranding records in this region, it appears that Hubb's beaked whale is most commonly encountered (Carretta *et al.* 2008, Moore and Barlow 2013). Insufficient sighting records exist off the U.S. west coast (Figure 1) to determine any possible spatial or seasonal patterns in the distribution of mesoplodont beaked whales.

Until methods of distinguishing these six species at-sea are developed, the management unit must be defined to include all *Mesoplodon* stocks in this region. However, in the future, species-level management is desirable, and a high priority should be placed on finding means to obtain species-specific abundance information. For the Marine Mammal Protection Act (MMPA) stock assessment reports, three *Mesoplodon* stocks are defined: 1) all *Mesoplodon* species off California, Oregon and Washington (this report), 2) *M. stejnegeri* in Alaskan waters, and 3) *M. densirostris* in Hawaiian waters.

### POPULATION SIZE

Although mesoplodont beaked whales have been sighted along the U.S. west coast on several line transect surveys utilizing both aerial and shipboard platforms, the rarity of sightings has historically precluded reliable population estimates. Early abundance estimates are imprecise and biased low by an unknown amount because of the large proportion of time this species spends submerged, and because the ship surveys before 1996 covered only California waters, and thus did not include animals off Oregon/Washington. Furthermore, survey data include a large number of unidentified beaked whale sightings that are probably either *Mesoplodon* sp. or Cuvier's beaked whales (*Ziphius cavirostris*). An abundance estimate of 1,024 (CV = 0.77) for all species of *Mesoplodon* beaked whales in the California Current was obtained based on combining data from the two most recent surveys (2005, 2008) conducted within 300 nmi of the coasts of California, Oregon and Washington (Forney 2007, Barlow and Forney 2007, Barlow 2010). This estimate was based in part on a correction factor to account for the



**Figure 1.** *Mesoplodon* beaked whale sightings based on shipboard surveys off California, Oregon and Washington, 1991-2008 (see Appendix 2 for data sources and information on timing and location of survey effort). Key: ● = *Mesoplodon* spp.; ▲ = identified *Mesoplodon densirostris*; ■ = identified *Mesoplodon carlhubbsi*. Dashed line represents the U.S. EEZ, thin lines indicate completed transect effort of all surveys combined.

proportion of animals on the survey trackline that were likely to be missed by observers (0.55), calculated from a model of beaked whale diving behavior, detection distances and searching behavior by the observers (Barlow 1999). Of the 5 sightings of *Mesoplodon* made during 2005-2008 surveys [all 5 sightings were made during the 2005 survey] two were identified to the ‘probable’ species level (one *Mesoplodon densirostris* and one *Mesoplodon carlhubbsi*). An estimate of Blainville’s beaked whale abundance (603, CV = 1.16) was based on this one probable sighting, while the Hubb’s beaked whale sighting was not recorded during standard survey effort, and thus there is no estimate of abundance. The abundance estimate for mesoplodont beaked whales of unknown species, based on the same 2005-2008 surveys was 421 (CV=0.88). A trend-based analysis of line-transect data from surveys conducted between 1991 and 2008 yielded new estimates of *Mesoplodon* species abundance (Moore and Barlow 2013). The new estimate accounts for the proportion of unidentified beaked whale sightings likely to be *Mesoplodon* beaked whales and uses a correction factor for missed animals adjusted to account for the fact that the proportion of animals on the trackline missed by observers increases in rough observing conditions. The trend-model analysis incorporates information from the entire 1991-2008 time series for each annual estimate of abundance, and given the strong evidence of a decreasing abundance trend over that time (Moore and Barlow 2013), the best estimate of abundance is represented by the model-averaged estimate for 2008. Based on this analysis, the best (50<sup>th</sup> percentile) estimate of abundance for all species of *Mesoplodon* species combined in 2008 in waters off California, Oregon and Washington is 694 (CV=0.65).

### Minimum Population Estimate

The minimum population estimate (defined as the log-normal 20th percentile of the abundance estimate) for mesoplodont beaked whales in California, Oregon, and Washington is 389 animals.

### Current Population Trend


There is strong evidence, based on line-transect survey data and the historical stranding record off the U.S. west coast, that the abundance of *Mesoplodon* beaked whales has recently declined in waters off California, Oregon and Washington (Moore and Barlow 2013, Figure 2). Statistical analysis of line-transect survey data from 1991 - 2008 indicates a 0.96 probability of decline during this period, with the mean annual rate of population change estimated to have been -7.0% per year (95% CRI: -16.7% to +1.0%). Patterns in the historical stranding record alone provide limited information about beaked whale abundance trends, but the stranding record appears generally consistent rather than at-odds with results of the line-transect survey analysis. Regional stranding networks along the Pacific coast of the U.S. and Canada originated during the 1980s, and beach coverage and reporting rates are thought to have increased throughout the 1990s and in to the early 2000s. Therefore, for a stable or increasing population, an overall increasing trend in stranding reports between the 1980s and 2000s would be expected. In contrast, reported strandings for *M. carlhubbsi* and *M. stejnegeri* in the California Current region have declined monotonically since the 1980s.

### CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

No information on current or maximum net productivity rates is available for mesoplodont beaked whales.

### POTENTIAL BIOLOGICAL REMOVAL

The potential biological removal (PBR) level for this stock is calculated as the minimum population size (389) times one half the default maximum net growth rate



**Figure 2.** Abundance and trend estimates for mesoplodont beaked whales in the California Current, 1991-2008 (Moore and Barlow 2013). For each year, the Bayesian posterior median (●), mean (x) and mode (\*) abundance estimates are shown, along with 90% CRIs.

for cetaceans (½ of 4%) times a recovery factor of 0.50 (for a species of unknown status with no known recent fishery mortality; Wade and Angliss 1997), resulting in a PBR of 3.9 mesoplodont beaked whales per year.

## HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

### Fishery Information

The California large mesh drift gillnet fishery has been the only fishery historically known to interact with *Mesoplodon* beaked whales in this region. Between 1990 and 1995, a total of eight *Mesoplodon* beaked whales (5 Hubb’s beaked whales (*Mesoplodon carlhubbsi*), one Stejneger’s beaked whale (*Mesoplodon stejnegeri*), and two unidentified whales of the genus *Mesoplodon* were observed entangled in approximately 3,300 sets (Julian and Beeson 1998, Carretta *et al.* 2008). Following the introduction of acoustic pingers into this fishery (Barlow and Cameron 2003), no beaked whales of any species have been observed entangled in over 4,000 observed sets (Carretta *et al.* 2008, Carretta and Enriquez 2009a, 2009b, 2010, 2012a, 2012b, Carretta and Barlow 2011). Mean annual takes in Table 1 are based on 2007-2011 data. This results in an average estimated annual mortality of zero mesoplodont beaked whales.

Gillnets have been documented to entangle marine mammals off Baja California (Sosa-Nishizaki *et al.* 1993), but no recent bycatch data from Mexico are available.

**Table 1.** Summary of available information on the incidental mortality and injury of *Mesoplodon* beaked whales (California/Oregon/Washington Stocks) in commercial fisheries that might take these species. Mean annual takes are based on 2007-2011 data unless noted otherwise.

Fishery Name	Data Type	Year	Percent Observer Coverage	Observed Mortality	Estimated Annual Mortality	Mean Annual Takes (CV in parentheses)
CA/OR thresher shark/swordfish drift gillnet fishery	observer	2007	16.4%	0	0	0
		2008	13.5%	0	0	
		2009	13.3%	0	0	
		2010	11.9%	0	0	
		2011	19.5%	0	0	
<b>Minimum total annual takes of <i>Mesoplodon</i> beaked whales</b>						0

### Other mortality

Anthropogenic sound sources, such as military sonar and seismic testing have been implicated in the mass strandings of beaked whales, including atypical events involving multiple beaked whale species (Simmonds and Lopez-Jurado 1991, Frantiz 1998, Anon. 2001, Jepson *et al.* 2003, Cox *et al.* 2006). While D’Amico *et al.* (2009) note that most mass strandings of beaked whales are unassociated with documented sonar activities, lethal or sub-lethal effects of such activities would rarely be documented, due to the remote nature of such activities and the low probability that an injured or dead beaked whale would strand. Filadelpho *et al.* (2009) reported statistically significant correlations between military sonar use and mass strandings of beaked whales in the Mediterranean and Caribbean Seas, but not in Japanese and Southern California waters, and hypothesized that regions with steep bathymetry adjacent to coastlines are more conducive to stranding events in the presence of sonar use. In Hawaiian waters, Faerber & Baird (2010) suggest that the probability of stranding is lower than in some other regions due to nearshore currents carrying animals away from beaches, and that stranded animals are less likely to be detected due to low human population density near many of Hawaii’s beaches. Actual and simulated sonar are known to interrupt the foraging dives and echolocation activities of tagged beaked whales (Tyack *et al.* 2011, DeRuiter *et al.* 2013). Cuvier’s beaked whales tagged and tracked during simulated mid-frequency sonar exposure showed avoidance reactions, including prolonged diving, cessation of echolocation click production associated with foraging, and directional travel away from the simulated sonar source (DeRuiter *et al.* 2013). Blainville’s beaked whale presence was monitored on hydrophone arrays before, during, and after sonar activities on a Caribbean military range, with evidence of avoidance behavior: whales were detected throughout the range prior to sonar exposure, not detected in the center of the range coincident with highest sonar use, and gradually returned to the range center after the cessation of sonar activity (Tyack *et al.* 2011). Fernández *et al.* (2013) report that there have been no mass strandings of beaked

whales in the Canary Islands following a 2004 ban on sonar activities in that region. The absence of beaked whale bycatch in California drift gillnets following the introduction of acoustic pingers into the fishery implies additional sensitivity of beaked whales to anthropogenic sound (Carretta et al. 2008, Carretta and Barlow 2011).

#### STATUS OF STOCKS

The status of mesoplodont beaked whales in California, Oregon and Washington waters relative to OSP is not known, but evidence suggests a high likelihood of population decline in the California Current since the early 1990s, at a mean rate of  $-7.0\%$  per year, which corresponds to trend-fitted abundance levels in 2008 (most recent survey) being at approximately 30% of 1991 levels. Moore and Barlow (2013) ruled out bycatch as a cause of the decline in mesoplodont beaked whale abundance and suggest that impacts from anthropogenic sound such as naval sonar and deepwater ecosystem changes within the California Current are plausible hypotheses warranting further investigation. None of the six species is listed as "threatened" or "endangered" under the Endangered Species Act, but given the long-term decline in mesoplodont beaked whale abundance in the California Current reported by Moore and Barlow (2013), these stocks are considered strategic. The degree of decline (trend-fitted 2008 abundance at approximately 30% of 1991 levels) also suggests that these stocks are likely well below their carrying capacity and may be depleted. The average annual known human-caused fishery mortality between 2007 and 2011 is zero. It is likely that the difficulty in identifying these animals in the field will remain a critical obstacle to obtaining species-specific abundance estimates and stock assessments in the future. The impacts of anthropogenic sound on beaked whales remains a concern (Barlow and Gisiner 2006, Cox et al. 2006, Hildebrand et al. 2005, Weilgart 2007).

#### REFERENCES

- Anon. 2001. Joint Interim Report, Bahamas Marine Mammal Stranding Event of 15\_16 March 2000. Available from NOAA, NMFS, Office of Protected Resources, Silver Spring, MD.
- Barlow, J. 2010. Cetacean abundance in the California Current from a 2008 ship-based line-transect survey. NOAA Technical Memorandum, NMFS, NOAA-TM-NMFS-SWFSC-456.
- Barlow, J. 1999. Trackline detection probability for long-diving whales. p. 209-224 *In*: G. W. Garner, S. C. Amstrup, J. L. Laake, B. F. J. Manly, L. L. McDonald, and D. G. Robertson (eds.) Marine Mammal Survey and Assessment Methods. A. A. Balkema, Rotterdam. 287 pp.
- Barlow, J. and G. A. Cameron. 2003. Field experiments show that acoustic pingers reduce marine mammal bycatch in the California drift gillnet fishery. *Marine Mammal Science* 19(2):265-283.
- Barlow, J. and K.A. Forney. 2007. Abundance and population density of cetaceans in the California Current ecosystem. *Fishery Bulletin* 105:509-526.
- Barlow, J. and R. Gisiner. 2006. Mitigating, monitoring, and assessing the effects of anthropogenic sound on beaked whales. *J. Cet. Res. Manage.* 7(3):239-249.
- Carretta, J.V., J. Barlow, and L. Enriquez. 2008. Acoustic pingers eliminate beaked whale bycatch in a gillnet fishery. *Marine Mammal Science* 24(4): 956-961.
- Carretta, J.V. and L. Enriquez. 2009a. Marine mammal and seabird bycatch observed in California commercial fisheries in 2007. Administrative Report LJ-09-01, available from Southwest Fisheries Science Center, 3333 North Torrey Pines Rd., La Jolla, CA 92037. 12 p.
- Carretta, J.V. and L. Enriquez. 2009b. Marine mammal bycatch observed in the California/Oregon swordfish and thresher shark drift gillnet fishery in 2008. Administrative Report LJ-09-03, available from Southwest Fisheries Science Center, 3333 North Torrey Pines Rd., La Jolla, CA 92037. 10 p.
- Carretta, J.V. and L. Enriquez. 2010. Marine Mammal and Sea Turtle Bycatch in the California/Oregon Swordfish and Thresher Shark Drift Gillnet Fishery in 2009. Southwest Fisheries Science Center, NOAA Fisheries Administrative Report LJ-10-03. 11p.
- Carretta, J.V. and L. Enriquez. 2012b. Marine mammal and seabird bycatch in California gillnet fisheries in 2011. NOAA Technical Memorandum NOAA-TM-NMFS-SWFSC-504. 14 p.
- Carretta, J.V. and L. Enriquez. 2012a. Marine mammal and seabird bycatch in California gillnet fisheries in 2010. Southwest Fisheries Science Center Administrative Report LJ-12-01. 14p.
- Carretta, J.V. and J. Barlow. 2011. Long-term effectiveness, failure rates, and "dinner bell" properties of acoustic pingers in a gillnet fishery. *Marine Technology Society Journal* 45:7-19.

- Cox, T. M., T. J. Ragen, A. J. Read, E. Vos, R. W. Baird, K. Balcomb, J. Barlow, J. Caldwell, T. Cranford, L. Crum, A. D'Amico, G. D'spain, A. Fernandez, J. Finneran, R. Gentry, W. Gerth, F. Gulland, J. Hildebrand, D. Houser, T. Hullar, P. D. Jepson, D. Ketten, C. D. Macleod, P. Miller, S. Moore, D. C. Mountain, D. Palka, P. Ponganis, S. Rommel, T. Rowles, B. Taylor, P. Tyack, D. Wartzok, R. Gisiner, J. Mead and L. Benner. 2006. Understanding the impacts of anthropogenic Sound on beaked whales. *Journal of cetacean research and management* 7:177–187.
- D'Amico A., Gisiner R.C., Ketten D.R., Hammock J.A., Johnson C., et al. 2009. Beaked whale strandings and naval exercises. *Aquat. Mamm.* 34: 452–472.
- Dalebout, M. L., J. G. Mead, C. Scott Baker, A. N. Baker and A. L. van Helden. (2002). A new species of beaked whale *Mesoplodon perrini* sp. n. (Cetacea: Ziphiidae) discovered through phylogenetic analyses of mitochondrial DNA sequences. *Marine Mammal Science* 18:577-608.
- DeRuiter, S.L., Southall B.L., Calambokidis J., Zimmer W.M.X., Sadykova D., Falcone E.A., Friedlaender A.S., Joseph J.E., Moretti D., Schorr G.S., Thomas L., Tyack P.L. 2013. First direct measurements of behavioural responses by Cuvier's beaked whales to mid-frequency active sonar. *Biol Lett* 9: 20130223. <http://dx.doi.org/10.1098/rsbl.2013.0223>
- Fernández, A., Arbelo, M. and Martín, V. 2013. No mass strandings since sonar ban. *Nature* 497:317.
- Filadelfo R., Mintz J., Michlovich E., D'Amico A., Tyack P.L. 2009. Correlating military sonar use with beaked whale mass strandings: what do the historical data show? *Aquat Mamm* 34: 435–444.
- Frantzis, A. 1998. Does acoustic testing strand whales? *Nature* 392(5):29.
- Forney, K.A. 2007. Preliminary estimates of cetacean abundance along the U.S. west coast and within four National Marine Sanctuaries during 2005. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-SWFSC-406. 27p.
- Henshaw, M.D., R.G. LeDuc, S.J. Chivers, and A.E. Dizon. 1997. Identifying beaked whales (Family Ziphiidae) using mtDNA sequences. *Marine Mammal Science* 13(3):487-495.
- Hildebrand, J.A. 2005. Impacts of anthropogenic sound. In: Reynolds III JE, Perrin WF, Reeves RR, Montgomery S, Ragen TJ, editors. *Marine mammal research: conservation beyond crisis*. Baltimore: Johns Hopkins University. pp. 101–123.
- Jepson, P.D., Arbelo, M., Deaville, R., Patterson, I.A.P., Castro, P., Baker, J.R., Degollada, E., Ross, H.M., Herraes, P., Pocknell, A.M., Rodriguez, F., Howiell, F.E., Espinosa, A., Reid, R.J., Jaber, J.R., Martin, V., Cunningham, A.A. and Fernández, A. 2003. Gas-bubble lesions in stranded animals: Was sonar responsible for a spate of whale deaths after an Atlantic military exercise? *Nature* 425(6958):575-76.
- Julian, F. and M. Beeson. 1998. Estimates of mammal, turtle and bird mortality for two California gillnet fisheries: 1990-1995. *Fishery Bulletin* 96:271-284.
- Mead, J. G. 1989. Beaked whales of the genus *Mesoplodon*. In: Ridgway, S. H. and Harrison, R. (eds.), *Handbook of Marine Mammals*, Vol. 4., p. 349-430. Academic Press Limited.
- MacLeod C.D., Perrin W.F., Pitman R., Barlow J., Ballance L., D'Amico, A., Gerrodette, T., Joyce, G., Mullin, K.D., Palka, D.L., and Waring, G.T. 2006. Known and inferred distributions of beaked whale species (Cetacea: Ziphiidae). *Journal of Cetacean Research and Management* 7:271–286.
- Moore, J.E., Barlow J.P. 2013. Declining abundance of beaked whales (family Ziphiidae) in the California Current large marine ecosystem. *PLoS ONE* 8(1):e52770. doi:10.1371/journal.pone.0052770
- National Marine Fisheries Service, Southwest Fisheries Science Center. Unpublished data.
- Richardson, W. J., C. R. Greene, Jr., C. I. Malme, and D. H. Thompson. 1995. *Marine Mammals and Noise*. Academic Press, San Diego. 576 p.
- Simmonds, M.P., and Lopez-Jurado, L.F. 1991. Whales and the military. *Nature (London)*, 351:448. doi:10.1038/351448a0.
- Sosa-Nishizaki, O., R. De la Rosa-Pacheco, R. Castro-Longoria, M. Grijalva Chon, and J. De la Rosa Velez. 1993. Estudio biológico pesquero del pez (*Xiphias gladius*) y otras especies de picudos (marlins y pez vela). Rep. Int. CICESE, CTECT9306.
- Tyack P.L., Zimmer W.M.X., Moretti D., Southall B.L., Claridge D.E., et al. (2011) Beaked Whales Respond to Simulated and Actual Navy Sonar. *PLoS ONE* 6(3):e17009. doi:10.1371/journal.pone.0017009.
- Wade, P. R. and R. P. Angliss. 1997. Guidelines for Assessing Marine Mammal Stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U. S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12. 93 pp.

Weilgart, L.S. 2007. The impacts of anthropogenic ocean noise on cetaceans and implications for management. *Canadian Journal of Zoology* 85:1091-1116.