

KILLER WHALE (*Orcinus orca*): AT1 Transient Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

Killer whales have been observed in all oceans and seas of the world (Leatherwood and Dahlheim 1978). Although reported from tropical and offshore waters, killer whales occur at higher densities in colder and more productive waters of both hemispheres, with the greatest densities found at high latitudes (Mitchell 1975, Leatherwood and Dahlheim 1978, Forney and Wade 2006). Killer whales are found throughout the North Pacific Ocean. Along the west coast of North America, seasonal and year-round occurrence of killer whales has been noted along the entire Alaska coast (Braham and Dahlheim 1982), in British Columbia and Washington inland waterways (Bigg et al. 1990), and along the outer coasts of Washington, Oregon, and California (Green et al. 1992; Barlow 1995, 1997; Forney et al. 1995). Killer whales from these areas have been labeled as “resident,” “transient,” and “offshore” type killer whales (Bigg et al. 1990, Ford et al. 2000, Dahlheim et al. 2008) based on aspects of morphology, ecology, genetics, and behavior (Ford and Fisher 1982; Baird and Stacey 1988; Baird et al. 1992; Hoelzel et al. 1998, 2002; Barrett-Lennard 2000; Dahlheim et al. 2008). Through examination of photographs of recognizable individuals and pods, movements of whales between geographical areas have been documented. For example, whales identified in Prince William Sound have been observed near Kodiak Island (Matkin et al. 1999) and whales identified in Southeast Alaska have been observed in Prince William Sound, British Columbia, and Puget Sound (Leatherwood et al. 1990, Dahlheim et al. 1997). Movements of killer whales between the waters of Southeast Alaska and central California have also been documented (Goley and Straley 1994, Black et al. 1997, Dahlheim and White 2010).

Several studies provide evidence that the resident, offshore, and transient ecotypes are genetically distinct in both mtDNA and nuclear DNA (Hoelzel and Dover 1991; Hoelzel et al. 1998, 2002; Barrett-Lennard 2000). Genetic differences have also been found between populations within the transient and resident ecotypes (Hoelzel et al. 1998, 2002; Barrett-Lennard 2000). A recent global genetic study of killer whales using the entire mitochondrial genome found that some killer whale ecotypes represent deeply divergent evolutionary lineages and warrant elevation to species or subspecies status (Morin et al. 2010). In particular, estimates from mitogenome sequence data indicate that transient killer whales diverged from all other killer whale lineages ~700,000 years ago. In light of these differences, the Society for Marine Mammalogy’s Committee on Taxonomy currently recognizes the resident and transient North Pacific ecotypes as un-named *Orcinus orca* subspecies (Committee on Taxonomy 2017). In recognition of its status as an un-named subspecies or species, some researchers now refer to transient-type killer whales as Bigg’s killer whales (e.g., Ford 2011, Riesch et al. 2012), in tribute to the late Dr. Michael Bigg.

The first studies of transient killer whales in Alaska were conducted in Southeast Alaska and in the Gulf of Alaska (from Prince William Sound, through the Kenai Fjords, and around Kodiak Island). In the Gulf of Alaska, Matkin et al. (1999) described two genetically distinct populations of transients which were never found in association with one another, the so-called “Gulf of Alaska” transients and “AT1” transients. In the past, neither of these populations were known to associate with the population of transient killer whales that ranged from California to Southeast Alaska, which are described as the West Coast Transient stock. Gulf of Alaska transients are documented throughout the Gulf of Alaska, including occasional sightings in Prince William Sound. AT1 transients have been seen only in Prince William Sound and in the Kenai Fjords region, and are therefore partially sympatric with Gulf of Alaska transients. In addition, recent data have identified 14 out of 217 transients on the outer coast of




Figure 1. Approximate distribution of transient killer whales in the eastern North Pacific (shaded areas). The distribution of resident and transient killer whale stocks in the eastern North Pacific largely overlap (see text).

Southeast Alaska and British Columbia as Gulf of Alaska transients and in one encounter they were observed mixing with West Coast Transients (Matkin et al. 2012, Ford et al. 2013). Transients within the Gulf of Alaska population have been found to have two mtDNA haplotypes, neither of which is found in the West Coast or AT1 populations. Members of the AT1 population share a single mtDNA haplotype. Transient killer whales from the West Coast population have been found to share a single mtDNA haplotype that is not found in the other populations. Additionally, all three populations have been found to have significant differences in nuclear (microsatellite) DNA (Barrett-Lennard 2000). Acoustic differences have been found as well; Saulitis et al. (2005) described acoustic differences between Gulf of Alaska transients and AT1 transients. For these reasons, the Gulf of Alaska transients are considered part of a population that is discrete from the AT1 population, and both of these populations are considered discrete from the West Coast Transients.

Biopsy samples from the eastern Aleutians and the south side of the west end of the Alaska Peninsula have produced the same haplotypes as killer whales in the northern Gulf of Alaska; however, nuclear DNA analysis strongly suggests they belong to a separate population (Parsons et al. 2013). The geographic distribution of mtDNA haplotypes revealed samples from the central Aleutian Islands and Bering Sea with haplotypes not found in Gulf of Alaska transients, suggesting additional population structure in western Alaska. At this time, transient-type killer whales from the Aleutian Islands and Bering Sea are considered to be part of a single population that includes Gulf of Alaska transients. Killer whales observed in the northern Bering Sea and Beaufort Sea have physical characteristics of transient-type whales, but little is known about these whales. AT1 haplotype whales are also present west of the Aleutian Islands and into the Bering Sea; however, nuclear DNA analysis indicates these animals are not part of the AT1 transient population in the Gulf of Alaska (L. Barrett-Lennard, Vancouver Aquarium, pers. comm., 21 March 2014).

In summary, within the transient ecotype, association data (Ford et al. 1994, Ford and Ellis 1999, Matkin et al. 1999), acoustic data (Ford and Ellis 1999, Saulitis et al. 2005), and genetic data (Hoelzel et al. 1998, 2002; Barrett-Lennard 2000) confirm that at least three communities of transient whales exist and represent three discrete populations: 1) Gulf of Alaska, Aleutian Islands, and Bering Sea transients, 2) AT1 transients, and 3) West Coast transients.

Based on data regarding association patterns, acoustics, movements, and genetic differences, eight killer whale stocks are now recognized within the Pacific U.S. Exclusive Economic Zone: 1) the Alaska Resident stock - occurring from Southeast Alaska to the Aleutian Islands and Bering Sea, 2) the Northern Resident stock - occurring from Washington State through part of Southeast Alaska, 3) the Southern Resident stock - occurring mainly within the inland waters of Washington State and southern British Columbia, but also in coastal waters from Southeast Alaska through California, 4) the Gulf of Alaska, Aleutian Islands, and Bering Sea Transient stock - occurring mainly from Prince William Sound through the Aleutian Islands and Bering Sea, 5) the AT1 Transient stock - occurring in Alaska from Prince William Sound through the Kenai Fjords (Fig. 1), 6) the West Coast Transient stock - occurring from California through Southeast Alaska, 7) the Offshore stock - occurring from California through Alaska, and 8) the Hawaiian stock. Transient whales in Canadian waters are considered part of the West Coast Transient stock. The Hawaiian and Offshore stocks are reported separately in the Stock Assessment Reports for the U.S. Pacific Region.

AT1 killer whales were first identified as a separate, cohesive group in 1984, when 22 transient-type whales were documented in Prince William Sound (Leatherwood et al. 1984, Heise et al. 1991), though individual whales from the group had been photographed as early as 1978 (von Ziegesar et al. 1986). Once the North Gulf Oceanic Society began consistent annual research effort in Prince William Sound, AT1 killer whales were resighted frequently. In fact, AT1 killer whales were found to be some of the most frequently sighted killer whales in Prince William Sound (Matkin et al. 1993, 1994, 1999). Gulf of Alaska transients are seen less frequently in Prince William Sound, with periods of several years or more between resightings.

AT1 killer whales have never been seen in association with sympatric resident killer whale pods or with Gulf of Alaska transients (Matkin et al. 1999, 2012), are genetically and acoustically distinct from other transient killer whales in the North Pacific (Barrett-Lennard 2000, Saulitis et al. 2005), and appear to have a more limited range than other transients. Their approximately 200-mile known range includes only Prince William Sound and Kenai Fjords and adjacent offshore waters (Matkin et al. 1999, 2012).

POPULATION SIZE

Using photographic-identification, all 22 individuals in the AT1 Transient population were censused for the first time in 1984 (Leatherwood et al. 1984). All 22 AT1 killer whales were seen annually or biannually from 1984 to 1988 (Matkin et al. 1999, 2003). The *Exxon Valdez* oil spill occurred in spring of 1989. Nine individuals from the AT1 group have been missing since 1990 (last seen in 1989), and two have been missing since 1992 (last seen in

1990 and 1991). Three of the missing AT1 killer whales (AT5, AT7, and AT8) were seen near the leaking *Exxon Valdez* shortly after the spill (Matkin et al. 1993, 1994, 2008). Two whales were found dead, stranded in 1989-1990, both genetically assigned to the AT1 population and one visually recognized as AT19, one of the missing nine (Matkin et al. 1994, 2008; Heise et al. 2003). The second unidentified whale was most likely one of the other missing AT1 whales. Additional mortalities of four older males include whales AT1 found stranded in 2000, AT13 and AT17 missing in 2002 (one of which was thought to be the carcass from the AT1 population that was found in 2002), and AT14 missing in 2003. A stranded whale found in 2003, genetically assigned to the AT1 population, was probably AT14 but could also have been AT13 (Matkin et al. 2008). No births have occurred in this population since 1984 and none of the missing whales have been seen since 2003 and are presumed dead. There is an extremely small probability (0.4%) that AT1 killer whales that are missing for 3 years or more are still alive (Matkin et al. 2008). No AT1 killer whale missing for at least 4 years has ever been resighted, and all 15 missing whales are presumed dead (Matkin et al. 2008). In 2016, six of the seven whales were photographed by researchers from the North Gulf Oceanic Society (birth year is estimated for whales born before 1983, as described in Matkin et al. 1999): AT2 (female, born ≤ 1969), AT3 (male, born 1984), AT4 (female, born ≤ 1974), AT9 (female, born ≤ 1965), AT10 (male, born 1980), and AT18 (female, born ≤ 1974). Researcher Craig Matkin believes he saw AT6 (male, born 1976) in 2016 but did not obtain a photograph. Therefore, the population estimate as of the summer of 2016 remains at seven whales (C. Matkin, North Gulf Oceanic Society, pers. comm., 27 September 2016). There has been no recruitment in this population since 1984 (Matkin et al. 2012).

Minimum Population Estimate

The abundance estimate of killer whales is a direct count of individually identifiable animals. Only 11 whales were seen between 1990 and 1999. Since then, four of those whales have not been seen for four or more consecutive years, so the minimum population estimate (N_{MIN}) is seven whales (Matkin et al. 2008; C. Matkin, pers. comm., 27 September 2016). Fourteen years of annual effort have failed to discover any whales that had not been seen previously, so there is no reason to believe there are additional whales in the population. Therefore, this N_{MIN} is the total population size.

Current Population Trend

The population counts have declined from a level of 22 whales in 1989 to 7 whales in 2015, a decline of 68%. Most of the mortality apparently occurred in 1989-1990.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

A reliable estimate of the maximum net productivity rate is unavailable for this stock of killer whales. Studies of resident killer whale pods in the Pacific Northwest resulted in estimated population growth rates of 2.9% and 2.5% over the period from 1973 to 1987 (Olesiuk et al. 1990, Brault and Caswell 1993). Until additional stock-specific data become available, the cetacean maximum theoretical net productivity rate (R_{MAX}) of 4% will be used for this stock (Wade and Angliss 1997).

POTENTIAL BIOLOGICAL REMOVAL

Potential biological removal (PBR) is defined as the product of the minimum population estimate, one-half the maximum theoretical net productivity rate, and a recovery factor: $\text{PBR} = N_{\text{MIN}} \times 0.5R_{\text{MAX}} \times F_R$. The recovery factor (F_R) for this stock is 0.1, as the stock is considered depleted under the Marine Mammal Protection Act (MMPA) and there has been no recruitment into the stock since 1984. Thus, for the AT1 Transient killer whale stock, $\text{PBR} = 0$ whales ($7 \times 0.02 \times 0.1$).

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Detailed information for each human-caused mortality, serious injury, and non-serious injury reported for NMFS-managed Alaska marine mammals in 2011-2015 is listed, by marine mammal stock, in Helker et al. (2017); however, only the mortality and serious injury data are included in the Stock Assessment Reports. No human-caused mortality or serious injury of AT1 Transient killer whales was reported in 2011-2015. Potential threats most likely to result in direct human-caused mortality or serious injury of this stock include ship strikes and oil spills (most of the mortality in this stock occurred in 1989-1990, following the *Exxon Valdez* oil spill).

Fisheries Information

Detailed information on U.S. commercial fisheries in Alaska waters (including observer programs, observer coverage, and observed incidental takes of marine mammals) is presented in Appendices 3-6 of the Alaska Stock Assessment Reports.

The known range of the AT1 Transient stock is limited to waters of Prince William Sound and Kenai Fjords. There are no federally-managed commercial fisheries in this area. State-managed commercial fisheries prosecuted within the range of this stock, such as the Prince William Sound salmon set and drift gillnet fisheries and various herring fisheries, are not known to cause incidental mortality or serious injury of AT1 killer whales. Several subsistence fisheries (salmon, halibut, non-salmon finfish, and shellfish) also occur within this area, and no incidental mortality or serious injury of AT1 killer whales has been reported for these fisheries.

Alaska Native Subsistence/Harvest Information

There are no reports of a subsistence harvest of killer whales in Alaska or Canada.

Other Mortality

Collisions with boats may be an occasional source of mortality or serious injury of killer whales. For example, a killer whale struck the propeller of a vessel in the Bering Sea/Aleutian Islands rockfish trawl fishery in 2010; however, this mortality did not involve a whale from the AT1 Transient stock. There has been no known mortality or serious injury of AT1 killer whales due to ship strikes. Most of the mortality occurred from 1989 to 1990 following the *Exxon Valdez* oil spill.

STATUS OF STOCK

The AT1 Transient stock of killer whales is below its Optimum Sustainable Population and designated as depleted under the MMPA (69 FR 31321, June 3, 2004); therefore, it is classified as a strategic stock. The AT1 Transient stock is not listed as threatened or endangered under the Endangered Species Act. Based on currently available data, the estimated mean annual mortality and serious injury rate due to U.S. commercial fisheries (0) does not exceed 10% of the PBR (0) and, therefore, can be considered insignificant and approaching zero mortality and serious injury rate. At least 11 animals were alive in 1998, but it appears that only 7 individuals remain alive. The AT1 killer whale group has been reduced to 32% (7/22) of its 1984 level. Since no births have occurred in the past 30 years, it is unlikely that this stock will recover.

There are few uncertainties in the assessment of the AT1 Transient stock of killer whales. Individual whales can be counted annually and the stock has been declining slowly since a dramatic reduction in the stock occurred immediately after the *Exxon Valdez* oil spill.

CITATIONS

- Baird, R. W., and P. J. Stacey. 1988. Variation in saddle patch pigmentation in populations of killer whales (*Orcinus orca*) from British Columbia, Alaska, and Washington State. *Can. J. Zool.* 66(11):2582-2585.
- Baird, R. W., P. A. Abrams, and L. M. Dill. 1992. Possible indirect interactions between transient and resident killer whales: implications for the evolution of foraging specializations in the genus *Orcinus*. *Oecologia* 89:125-132.
- Barlow, J. 1995. The abundance of cetaceans in California waters. Part I: Ship surveys in summer and fall of 1991. *Fish. Bull.*, U.S. 93:1-14.
- Barlow, J. 1997. Preliminary estimates of cetacean abundance off California, Oregon and Washington based on a 1996 ship survey and comparisons of passing and closing modes. Southwest Fisheries Science Center Administrative Report LJ-97-11. 25 p. Available from SWFSC, NMFS, 8901 La Jolla Shores Drive, La Jolla, CA 92037.
- Barrett-Lennard, L. G. 2000. Population structure and mating patterns of killer whales as revealed by DNA analysis. Ph.D. Dissertation, University of British Columbia, Vancouver, BC, Canada. 97 p.
- Bigg, M. A., P. F. Olesiuk, G. M. Ellis, J. K. B. Ford, and K. C. Balcomb III. 1990. Social organization and genealogy of resident killer whales (*Orcinus orca*) in the coastal waters of British Columbia and Washington State, p. 386-406. *In* P. S. Hammond, S. A. Mizroch, and G. P. Donovan (eds.), *Individual Recognition of Cetaceans: Use of Photo-identification and Other Techniques to Estimate Population Parameters*. *Rep. Int. Whal. Comm. Special Issue* 12.
- Black, N. A., A. Schulman-Janiger, R. L. Ternullo, and M. Guerrero-Ruiz. 1997. Killer whales of California and western Mexico: a catalog of photo-identified individuals. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-SWFSC-247, 174 p.

- Braham, H. W., and M. E. Dahlheim. 1982. Killer whales in Alaska documented in the Platforms of Opportunity Program. *Rep. Int. Whal. Comm.* 32:643-646.
- Brault, S., and H. Caswell. 1993. Pod-specific demography of killer whales (*Orcinus orca*). *Ecology* 74(5):1444-1454.
- Committee on Taxonomy. 2017. List of marine mammal species and subspecies. Society for Marine Mammalogy. Available online: www.marinemammalscience.org. Accessed December 2017.
- Dahlheim, M. E., and P. A. White. 2010. Ecological aspects of transient killer whales (*Orcinus orca*) as predators in southeastern Alaska. *Wildl. Biol.* 16:308-322.
- Dahlheim, M. E., D. Ellifrit, and J. Swenson. 1997. Killer Whales of Southeast Alaska: A Catalogue of Photoidentified Individuals. Day Moon Press, Seattle, WA. 82 p. + appendices.
- Dahlheim, M. E., A. Schulman-Janiger, N. Black, R. Ternullo, D. Ellifrit, and K. C. Balcomb. 2008. Eastern temperate North Pacific offshore killer whales (*Orcinus orca*): occurrence, movements, and insights into feeding ecology. *Mar. Mammal Sci.* 24:719-729.
- Ford, J. K. B. 2011. Killer whales of the Pacific Northwest coast: from pest to paragon. *Whale Watcher* 40(1):15-23.
- Ford, J. K. B., and G. M. Ellis. 1999. Transients: Mammal-Hunting Killer Whales of British Columbia, Washington, and Southeastern Alaska. University of British Columbia Press, Vancouver, BC. 96 p.
- Ford, J. K. B., and H. D. Fisher. 1982. Killer whale (*Orcinus orca*) dialects as an indicator of stocks in British Columbia. *Rep. Int. Whal. Comm.* 32:671-679.
- Ford, J. K. B., G. M. Ellis, and K. C. Balcomb. 1994. Killer Whales: The Natural History and Genealogy of *Orcinus orca* in British Columbia and Washington State. University of British Columbia Press, Vancouver, BC, and University of Washington Press, Seattle. 102 p.
- Ford, J. K. B., G. M. Ellis, and K. C. Balcomb. 2000. Killer Whales: The Natural History and Genealogy of *Orcinus orca* in British Columbia and Washington State. Second edition. University of British Columbia Press, Vancouver, BC, Canada. 104 p.
- Ford, J. K. B., E. H. Stredulinsky, J. R. Towers, and G. M. Ellis. 2013. Information in support of the identification of critical habitat for transient killer whales (*Orcinus orca*) off the west coast of Canada. DFO Canadian Science Advisory Secretariat Research Document 2012/nnn.
- Forney, K. A., and P. R. Wade. 2006. World-wide abundance and density of killer whales, p. 145-162. *In* J. A. Estes, D. P. DeMaster, D. F. Doak, T. M. Williams, and R. L. Brownell, Jr. (eds.), *Whales, Whaling, and Ocean Ecosystems*. University of California Press.
- Forney, K. A., J. Barlow, and J. V. Carretta. 1995. The abundance of cetaceans in California waters. Part II: Aerial surveys in winter and spring of 1991 and 1992. *Fish. Bull.*, U.S. 93:15-26.
- Goley, P. D., and J. M. Straley. 1994. Attack on gray whales (*Eschrichtius robustus*) in Monterey Bay, California, by killer whales (*Orcinus orca*) previously identified in Glacier Bay, Alaska. *Can. J. Zool.* 72:1528-1530.
- Green, G. A., J. J. Brueggeman, R. A. Grotefendt, C. E. Bowlby, M. L. Bonnell, and K. C. Balcomb. 1992. Cetacean distribution and abundance off Oregon and Washington, 1989-1990, p. 1-100. *In* J. J. Brueggeman (ed.), *Oregon and Washington marine mammal and seabird surveys. Final Report OCS Study MMS 91-0093*.
- Heise, K., G. Ellis, and C. Matkin. 1991. A Catalogue of Prince William Sound Killer Whales. North Gulf Oceanic Society, Homer, AK. 51 p.
- Heise, K., L. G. Barrett-Lennard, E. Saulitis, C. Matkin, and D. Bain. 2003. Examining the evidence for killer whale predation on Steller sea lions in British Columbia and Alaska. *Aquat. Mamm.* 29(3):325-334.
- Helker, V. T., M. M. Muto, K. Savage, S. Teerlink, L. A. Jemison, K. Wilkinson, and J. Jannot. 2017. Human-caused mortality and injury of NMFS-managed Alaska marine mammal stocks, 2011-2015. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-354, 112 p.
- Hoelzel, A. R., and G. A. Dover. 1991. Genetic differentiation between sympatric killer whale populations. *Heredity* 66:191-195.
- Hoelzel, A. R., M. E. Dahlheim, and S. J. Stern. 1998. Low genetic variation among killer whales (*Orcinus orca*) in the eastern North Pacific, and genetic differentiation between foraging specialists. *J. Hered.* 89:121-128.
- Hoelzel, A. R., A. Natoli, M. Dahlheim, C. Olavarria, R. Baird, and N. Black. 2002. Low worldwide genetic diversity in the killer whale (*Orcinus orca*): implications for demographic history. *Proc. R. Soc. Lond.* 269:1467-1473.
- Leatherwood, J. S., and M. E. Dahlheim. 1978. Worldwide distribution of pilot whales and killer whales. Naval Ocean Systems Center, Tech. Rep. 443:1-39.

- Leatherwood, S., A. E. Bowles, E. Krygier, J. D. Hall, and S. Ingell. 1984. Killer whales (*Orcinus orca*) in Southeast Alaska, Prince William Sound, and Shelikof Strait: a review of available information. Rep. Int. Whaling Comm. 34:521-530.
- Leatherwood, S., C. O. Matkin, J. D. Hall, and G. M. Ellis. 1990. Killer whales, *Orcinus orca*, photo-identified in Prince William Sound, Alaska 1976 to 1987. Can. Field Nat. 104:362-371.
- Matkin, C. O., M. E. Dahlheim, G. Ellis, and E. Saulitis. 1993. Vital rates and pod structure of resident killer whales following the *Exxon Valdez* oil spill, p. 303-307. In *Exxon Valdez Oil Spill Trustee Council, Exxon Valdez oil spill symposium abstract book*, February 2-5, 1993, Anchorage, Alaska.
- Matkin, C. O., G. M. Ellis, M. E. Dahlheim, and J. Zeh. 1994. Status of killer whales in Prince William Sound, 1985-1992, p. 141-162. In T. R. Loughlin (ed.), *Marine Mammals and the Exxon Valdez*. Academic Press, San Diego, CA.
- Matkin, C., G. Ellis, E. Saulitis, L. Barrett-Lennard, and D. Matkin. 1999. *Killer Whales of Southern Alaska*. North Gulf Oceanic Society, Homer, AK. 96 p.
- Matkin, C. O., G. M. Ellis, L. G. Barrett-Lennard, H. Yurk, E. L. Saulitis, D. Scheel, P. Olesiuk, and G. Ylitalo. 2003. Photographic and acoustic monitoring of killer whales in Prince William and Kenai Fjords, *Exxon Valdez Oil Spill Restoration Project Final Report*, Restoration Project 03012, North Gulf Oceanic Society, Homer, AK. 118 p.
- Matkin, C. O., E. L. Saulitis, G. M. Ellis, P. Olesiuk, and S. D. Rice. 2008. Ongoing population-level impacts on killer whales *Orcinus orca* following the 'Exxon Valdez' oil spill in Prince William Sound, Alaska. Mar. Ecol. Prog. Ser. 356:269-281.
- Matkin, C. O., J. W. Durban, E. L. Saulitis, R. D. Andrews, J. M. Straley, D. R. Matkin, and G. M. Ellis. 2012. Contrasting abundance and residency patterns of two sympatric populations of transient killer whales (*Orcinus orca*) in the northern Gulf of Alaska. Fish. Bull., U.S. 110:143-155.
- Mitchell, E. D. 1975. Report on the meeting on small cetaceans, Montreal, April 1-11, 1974. J. Fish. Res. Board Can. 32:914-916.
- Morin, P. A., F. I. Archer, A. D. Foote, J. Vilstrup, E. E. Allen, P. Wade, J. Durban, K. Parsons, R. Pitman, L. Li, P. Bouffard, S. C. A. Nielsen, M. Rasmussen, E. Willerslev, M. T. P. Gilbert, and T. Harkins. 2010. Complete mitochondrial genome phylogeographic analysis of killer whales (*Orcinus orca*) indicates multiple species. Genome Res. 20:908-916.
- Olesiuk, P. F., M. A. Bigg, and G. M. Ellis. 1990. Life history and population dynamics of resident killer whales (*Orcinus orca*) in the coastal waters of British Columbia and Washington State. Rep. Int. Whal. Comm. (Special Issue 12):209-242.
- Parsons, K. M., J. W. Durban, A. M. Burdin, V. N. Burkanov, R. L. Pitman, J. Barlow, L. G. Barrett-Lennard, R. G. LeDuc, K. M. Robertson, C. O. Matkin, and P. R. Wade. 2013. Geographic patterns of genetic differentiation among killer whales in the northern North Pacific. J. Hered. 104:737-754.
- Riesch, R., L. G. Barrett-Lennard, G. M. Ellis, J. K. B. Ford, and V. B. Deecke. 2012. Cultural traditions and the evolution of reproductive isolation: ecological speciation in killer whales? Biol. J. Linn. Soc. 106:1-17.
- Saulitis, E., C. O. Matkin, and F. H. Fay. 2005. Vocal repertoire and acoustic behavior of the isolated AT1 killer whale subpopulation in southern Alaska. Can. J. Zool. 83:1015-1029.
- von Ziegler, O., G. M. Ellis, C. O. Matkin, and B. Goodwin. 1986. Repeated sightings of identifiable killer whales (*Orcinus orca*) in Prince William Sound, Alaska, 1977-1983. Cetus 6(2):9-13.
- Wade, P. R., and R. Angliss. 1997. Guidelines for assessing marine mammal stocks: report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 p.