


## PACIFIC WHITE-SIDED DOLPHIN (*Lagenorhynchus obliquidens*): North Pacific Stock

### STOCK DEFINITION AND GEOGRAPHIC RANGE

The Pacific white-sided dolphin is found throughout the temperate North Pacific Ocean, north of the coasts of Japan and Baja California, Mexico. In the eastern North Pacific the species occurs from the southern Gulf of California, north to the Gulf of Alaska, west to Amchitka in the Aleutian Islands, and is rarely encountered in the southern Bering Sea. The species is common both on the high seas and along the continental margins, and animals are known to enter the inshore passes of Alaska, British Columbia, and Washington (Ferrero and Walker 1996).

The following information was considered in classifying Pacific white-sided dolphin stock structure based on the Dizon et al. (1992) phylogeographic approach: 1) Distributional data: geographic distribution is continuous; 2) Population response data: unknown; 3) Phenotypic data: two morphological forms are recognized (Walker et al. 1986, Chivers et al. 1993); and 4) Genotypic data: preliminary genetic analyses on 116 Pacific white-sided dolphins collected in four areas (Baja California, the U.S. west coast, British Columbia/Southeast Alaska, and offshore) do not support phylogeographic partitioning, though they are sufficiently differentiated to be treated as separate management units (Lux et al. 1997). This limited information is not sufficient to define stock structure throughout the North Pacific beyond the generalization that a northern form occurs north of about 33°N from southern California along the coast to Alaska and a southern form ranges from about 36°N southward along the coasts of California and Baja California, while the core of the population ranges across the North Pacific to Japan at latitudes south of 45°N. Data are lacking to determine whether this latter group might include animals from one or both of the coastal forms. Although the genetic data are unclear, management issues support the designation of two stocks; because the California and Oregon thresher shark/swordfish drift gillnet fishery (operating between 33°N and approximately 47°N) and, to a lesser extent, the groundfish and salmon fisheries in Alaska are known to interact with Pacific white-sided dolphins, two management stocks are recognized: 1) the California/Oregon/Washington stock, and 2) the North Pacific stock (Fig. 1). The California/Oregon/Washington stock is reported separately in the Stock Assessment Reports for the U.S. Pacific Region.



**Figure 1.** Approximate distribution of Pacific white-sided dolphins in the eastern North Pacific (dark shaded areas).

### POPULATION SIZE

The most complete population abundance estimate for Pacific white-sided dolphins was calculated from line-transect analyses applied to the 1987-1990 central North Pacific marine mammal sighting survey data (Buckland et al. 1993). The Buckland et al. (1993) abundance estimate, 931,000 (CV = 0.90) animals, more closely reflects a range-wide estimate rather than one that can be applied to either of the two management stocks off the west coast of North America. Furthermore, Buckland et al. (1993) suggested that Pacific white-sided dolphins show strong vessel attraction but that a correction factor was not available to apply to the estimate. While the Buckland et al. (1993) abundance estimate is not considered appropriate to apply to the management stock in Alaskan waters, the portion of the estimate derived from sightings north of 45°N in the Gulf of Alaska can be used as the population estimate for this area (26,880). For comparison, Hobbs and Lerczak (1993) estimated 15,200 (95% CI: 868-265,000) Pacific white-sided dolphins in the Gulf of Alaska based on a single sighting of 20 animals. Small cetacean aerial surveys in the Gulf of Alaska during 1997 sighted one group of 164 Pacific white-sided dolphins off Dixon entrance, while similar surveys in Bristol Bay in 1999 made 18 sightings of a school, or parts thereof, off Port Moller (R. Hobbs, NMFS-AFSC-NMML, pers. comm.).

### **Minimum Population Estimate**

Historically, the minimum population estimate ( $N_{\text{MIN}}$ ) for this stock was 26,880, based on the sum of abundance estimates for four separate  $5^\circ \times 5^\circ$  blocks north of  $45^\circ\text{N}$  ( $1,970 + 6,427 + 6,101 + 12,382 = 26,880$ ) from surveys conducted during 1987-1990, reported in Buckland et al. (1993). This was considered a minimum estimate because the abundance of animals in a fifth  $5^\circ \times 5^\circ$  block (53,885), which straddled the boundary of the two coastal management stocks, was not included in the estimate for the North Pacific stock and because much of the potential habitat for this stock was not surveyed between 1987 and 1990. However, because the abundance estimate is more than 8 years old, the current minimum population estimate for this stock is unknown.

### **Current Population Trend**

At present, there is no reliable information on trends in abundance for this stock of Pacific white-sided dolphins.

### **CURRENT AND MAXIMUM NET PRODUCTIVITY RATES**

A reliable estimate of the maximum net productivity rate is not currently available for the North Pacific stock of Pacific white-sided dolphins. Life-history analyses by Ferrero and Walker (1996) suggest a reproductive strategy consistent with the delphinid pattern on which the 4% cetacean maximum net productivity rate ( $R_{\text{MAX}}$ ) was based. Thus, it is recommended that the cetacean maximum net productivity rate ( $R_{\text{MAX}}$ ) of 4% be employed for this stock (Wade and Angliss 1997).

### **POTENTIAL BIOLOGICAL REMOVAL**

Under the 1994 reauthorized Marine Mammal Protection Act (MMPA), the potential biological removal (PBR) is defined as the product of the minimum population estimate, one-half the maximum theoretical net productivity rate, and a recovery factor:  $\text{PBR} = N_{\text{MIN}} \times 0.5R_{\text{MAX}} \times F_{\text{R}}$ . The recovery factor ( $F_{\text{R}}$ ) for this stock is 0.5, the value for cetacean stocks of unknown status (Wade and Angliss 1997). The estimate of abundance for Pacific white-sided dolphins is more than 8 years old; Wade and Angliss (1997) recommend that abundance estimates older than 8 years no longer be used to calculate a PBR level. In addition, there is no corroborating evidence from recent surveys in Alaska that provide abundance estimates for a portion of the stock's range or any indication of the current status of this stock. Thus, the PBR for this stock is undetermined (NMFS 2005).

### **ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY**

#### **New Serious Injury Guidelines**

NMFS updated its serious injury designation and reporting process, which uses guidance from previous serious injury workshops, expert opinion, and analysis of historic injury cases to develop new criteria for distinguishing serious from non-serious injury (Angliss and DeMaster 1998, Andersen et al. 2008, NOAA 2012). NMFS defines serious injury as an “*injury that is more likely than not to result in mortality.*” Injury determinations for stock assessments revised in 2013 or later incorporate the new serious injury guidelines, based on the most recent 5-year period for which data are available.

#### **Fisheries Information**

Between 1978 and 1991, mortality and serious injury of thousands of Pacific white-sided dolphins occurred annually incidental to high-seas fisheries for salmon and squid. However, these fisheries were closed in 1991 and no other large-scale fisheries have operated in the central North Pacific since 1991.

Detailed information (including observer programs, observer coverage, and observed incidental takes of marine mammals) for federally-managed and state-managed U.S. commercial fisheries in Alaska waters is presented in Appendices 3-6 of the Alaska Stock Assessment Reports.

Until 2003, there were six different federally-regulated commercial fisheries in Alaska that could have interacted with Pacific white-sided dolphins. These fisheries were monitored for incidental mortality and serious injury by fishery observers. As of 2003, changes in fishery definitions in the MMPA List of Fisheries have resulted in separating these 6 fisheries into 22 fisheries (69 FR 70094, 2 December 2004). This change does not represent a change in fishing effort but provides managers with better information on the component of each fishery that is responsible for the incidental mortality or serious injury of marine mammal stocks in Alaska. No mortality or serious injury of Pacific white-sided dolphins incidental to observed U.S. commercial fisheries was reported between 2009 and 2013 (Breiwick 2013; NMML, unpubl. data).

Note that no observers have been assigned to several of the gillnet fisheries that are known to interact with this stock, making the estimated mortality and serious injury rate unreliable. However, because the stock size is large, it is unlikely that unreported mortality and serious injury from those fisheries would be significant.

#### **Alaska Native Subsistence/Harvest Information**

There are no reports of subsistence takes of Pacific white-sided dolphins in Alaska.

#### **Other Mortality**

From 2009 to 2013, no human-caused mortality or serious injury of Pacific white-sided dolphins was reported to the NMFS Alaska Region stranding database (Helker et al. 2015).

#### **STATUS OF STOCK**

Pacific white-sided dolphins are not designated as “depleted” under the MMPA or listed as “threatened” or “endangered” under the Endangered Species Act. The North Pacific stock of Pacific white-sided dolphins is not classified as a strategic stock. Because the PBR for Pacific white-sided dolphins is undetermined, the level of human-caused mortality and serious injury relative to PBR is unknown and the level of annual U.S. commercial fishery-related mortality and serious injury that can be considered insignificant and approaching zero mortality and serious injury rate is unknown. Population trends and status of this stock relative to its Optimum Sustainable Population are currently unknown.

#### **HABITAT CONCERNS**

While the majority of Pacific white-sided dolphins are found throughout the North Pacific, there are also significant numbers found in shelf break and deeper nearshore areas. Thus, they are subject to a variety of habitat impacts. Of particular concern are nearshore areas, bays, channels, and inlets where some Pacific white-sided dolphins are vulnerable to physical modifications of nearshore habitats, resulting from urban and industrial development (including waste management and nonpoint source runoff), and noise (Linnenschmidt et al. 2013).

#### **CITATIONS**

- Andersen, M. S., K. A. Forney, T. V. N. Cole, T. Eagle, R. Angliss, K. Long, L. Barre, L. Van Atta, D. Borggaard, T. Rowles, B. Norberg, J. Whaley, and L. Engleby. 2008. Differentiating serious and non-serious injury of marine mammals: report of the Serious Injury Technical Workshop, 10-13 September 2007, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-39, 94 p.
- Angliss, R. P., and D. P. DeMaster. 1998. Differentiating serious and non-serious injury of marine mammals taken incidental to commercial fishing operations. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-13, 48 p.
- Breiwick, J. M. 2013. North Pacific marine mammal bycatch estimation methodology and results, 2007-2011. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-260, 40 p.
- Buckland, S. T., K. L. Cattanach, and R. C. Hobbs. 1993. Abundance estimates of Pacific white-sided dolphin, northern right whale dolphin, Dall’s porpoise and northern fur seal in the North Pacific, 1987/90. Pp. 387-407 *In* W. Shaw, R. L. Burgner, and J. Ito (eds.), *Biology, distribution and stock assessment of species caught in the high seas driftnet fisheries in the North Pacific Ocean*. International North Pacific Fisheries Commission Symposium; 4-6 November 1991, Tokyo, Japan.
- Chivers, S. J., K. M. Peltier, W. T. Norman, P. A. Akin, and J. Heyning. 1993. Population structure of cetaceans in California coastal waters. Paper SOCCS9 presented at the Status of California Cetacean Stocks Workshop, held in La Jolla, California, March 31-April 2, 1993. 49 pp.
- Dizon, A. E., C. Lockyer, W. F. Perrin, D. P. DeMaster, and J. Sisson. 1992. Rethinking the stock concept: a phylogeographic approach. *Conserv. Biol.* 6:24-36.
- Ferrero, R. C., and W. A. Walker. 1996. Age, growth, and reproductive patterns of the Pacific white-sided dolphin (*Lagenorhynchus obliquidens*) taken in high seas drift nets in the central North Pacific Ocean. *Can. J. Zool.* 74(9):1673-1687.
- Helker, V. T., B. M. Allen, and L. A. Jemison. 2015. Human-caused injury and mortality of NMFS-managed Alaska marine mammal stocks, 2009-2013. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-AFSC-300, 94 p.
- Hobbs, R. C., and J. A. Lerczak. 1993. Abundance of Pacific white-sided dolphin and Dall’s porpoise in Alaska estimated from sightings in the North Pacific Ocean and the Bering Sea during 1987-1991. 13 pp. Available from NMML, AFSC, NMFS, 7600 Sand Point Way NE, Seattle, WA 98115.

- Linnenschmidt, M., J. Teilmann, T. Akamatsu, R. Dietz, and L. A. Miller. 2013. Biosonar, dive, and foraging activity of satellite tracked harbor porpoises (*Phocoena phocoena*). *Mar. Mammal Sci.* 29(2):77-97.
- Lux, C. A., A. S. Costa, and A. E. Dizon. 1997. Mitochondrial DNA population structure of the Pacific white-sided dolphin. *Rep. Int. Whal. Comm.* 47:645-652.
- NMFS (National Marine Fisheries Service). 2005. Revisions to guidelines for assessing marine mammal stocks. 24 pp. Available online: <http://www.nmfs.noaa.gov/pr/pdfs/sars/gamms2005.pdf>. Accessed December 2015.
- NOAA. 2012. Federal Register 77:3233. National policy for distinguishing serious from non-serious injury of marine mammals. Available online: <http://www.nmfs.noaa.gov/op/pds/documents/02/238/02-238-01.pdf>. Accessed December 2015.
- Wade, P. R., and R. Angliss. 1997. Guidelines for assessing marine mammal stocks: report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 p.
- Walker, W. A., S. Leatherwood, K. R. Goodrich, W. F. Perrin, and R. K. Stroud. 1986. Geographical variation and biology of the Pacific white-sided dolphin, *Lagenorhynchus obliquidens*, in the north-eastern Pacific. Pp. 441-465 *In* M. M. Bryden and R. Harrison (eds.), *Research on Dolphins*. Clarendon Press, Oxford.