

HARBOR PORPOISE (*Phocoena phocoena phocoena*): Gulf of Maine/Bay of Fundy Stock

STOCK DEFINITION AND GEOGRAPHIC RANGE

This stock is found in U.S. and Canadian Atlantic waters. The distribution of harbor porpoises has been documented by sighting surveys, strandings and takes reported by NMFS observers in the Sea Sampling Programs. During summer (July to September), harbor porpoises are concentrated in the northern Gulf of Maine and southern Bay of Fundy region, generally in waters less than 150 m deep (Gaskin 1977; Kraus et al. 1983; Palka 1995), with a few sightings in the upper Bay of Fundy and on Georges Bank (Palka 2000). During fall (October–December) and spring (April–June), harbor porpoises are widely dispersed from New Jersey to Maine, with lower densities farther north and south. They are seen from the coastline to deep waters (>1800 m; Westgate et al. 1998), although the majority of the population is found over the continental shelf. During winter (January to March), intermediate densities of harbor porpoises can be found in waters off New Jersey to North Carolina, and lower densities are found in waters off New York to New Brunswick, Canada. Passive acoustic monitoring detected harbor porpoises regularly during the period January–May offshore of Maryland (Wingfield et al. 2017). There does not appear to be a temporally coordinated migration or a specific migratory route to and from the Bay of Fundy region. However, during the fall, several satellite-tagged harbor porpoises did favor the waters around the 92-m isobath, which is consistent with observations of high rates of incidental catches in this depth range (Read and Westgate 1997). There were two stranding records from Florida during the 1980s (Smithsonian strandings database) and one in 2003 (NE Regional Office/NMFS strandings and entanglement database).

Gaskin (1984, 1992) proposed that there were four separate populations in the western North Atlantic: the Gulf of Maine/Bay of Fundy, Gulf of St. Lawrence, Newfoundland, and Greenland populations. Analyses involving mtDNA (Wang et al. 1996; Rosel et al. 1999a; 1999b), organochlorine contaminants (Westgate et al. 1997; Westgate and Tolley 1999), heavy metals (Johnston 1995), and life history parameters (Read and Hohn 1995) support Gaskin's proposal. Genetic studies using mitochondrial DNA (Rosel et al. 1999a) and contaminant studies using total PCBs (Westgate and Tolley 1999) indicate that the Gulf of Maine/Bay of Fundy females were distinct from females from the other populations in the Northwest Atlantic. Gulf of Maine/Bay of Fundy males were distinct from Newfoundland and Greenland males, but not from Gulf of St. Lawrence males according to studies comparing mtDNA (Palka et al. 1996; Rosel et al. 1999a) and CHLORs, DDTs, PCBs and CHBs (Westgate and Tolley 1999). Nuclear microsatellite markers have also been




Figure 1. Distribution of harbor porpoises from NEFSC and SEFSC shipboard and aerial surveys during the summers of 1995, 1998, 1999, 2002, 2004, 2006, 2007, 2008, 2010, and 2011 and DFO's 2007 TNASS survey. Isobaths are the 100-m, 1000-m, and 4000-m depth contours.

applied to samples from these four populations, but this analysis failed to detect significant population sub-division in either sex (Rosel et al. 1999a). These patterns may be indicative of female philopatry coupled with dispersal of males. Both mitochondrial DNA and microsatellite analyses indicate that the Gulf of Maine/Bay of Fundy stock is not the sole contributor to the aggregation of porpoises found off the mid-Atlantic states during winter (Rosel et al. 1999a; Hiltunen 2006). Mixed-stock analyses using twelve microsatellite loci in both Bayesian and likelihood frameworks indicate that the Gulf of Maine/Bay of Fundy is the largest contributor (~60%), followed by Newfoundland (~25%) and then the Gulf of St. Lawrence (~12%), with Greenland making a small contribution (<3%). For Greenland, the lower confidence interval of the likelihood analysis includes zero. For the Bayesian analysis, the lower 2.5% posterior quantiles include zero for both Greenland and the Gulf of St. Lawrence. Intervals that reach zero provide the possibility that these populations contribute no animals to the mid-Atlantic aggregation.

This report follows Gaskin's hypothesis on harbor porpoise stock structure in the western North Atlantic, where the Gulf of Maine and Bay of Fundy harbor porpoises are recognized as a single management stock separate from harbor porpoise populations in the Gulf of St. Lawrence, Newfoundland, and Greenland. It is unlikely that the Gulf of Maine/Bay of Fundy harbor porpoise stock contains multiple demographically independent populations (Rosel et al. 1999a; Hiltunen 2006), but a comparison of samples from the Scotian shelf to the Gulf of Maine has not yet been made.

POPULATION SIZE

The best current abundance estimate of the Gulf of Maine/Bay of Fundy harbor porpoise stock is from the 2011 survey: 79,883 (CV=0.32). Key uncertainties include: 1) the surveyed area may not have covered the entire area of the stock's habitat at the appropriate time of the year, and 2) the current abundance estimate did not account for availability bias due to submergence of animals. Without a correction for availability bias, the abundance estimate is expected to be biased low. Since the dive times of harbor porpoises are relatively short (~ 4 minutes), it is expected the bias is not large.

Earlier abundance estimates

Please see Appendix IV for a summary of abundance estimates, including earlier estimates and survey descriptions. As recommended in the GAMMS II Workshop Report (Wade and Angliss 1997), estimates older than eight years are deemed unreliable for the determination of the current PBR.

Recent surveys and abundance estimates

An abundance estimate of 79,883 (CV=0.32) harbor porpoises was generated from a shipboard and aerial survey conducted during June–August 2011 (Palka 2012). The aerial portion that contributed to the abundance estimate covered 5,313 km of tracklines that were over waters north of New Jersey from the coastline to the 100-m depth contour through the U.S. and Canadian Gulf of Maine and up to and including the lower Bay of Fundy. The shipboard portion covered 3,107 km of tracklines that were in waters offshore of central Virginia to Massachusetts (waters that were deeper than the 100-m depth contour out to beyond the U.S. EEZ). Both sighting platforms used a double-platform team data-collection procedure, which allows estimation of abundance corrected for perception bias of the detected species (Laake and Borchers 2004). Estimation of the abundance was based on the independent-observer approach assuming point independence (Laake and Borchers 2004) and calculated using the mark-recapture distance sampling option in the computer program Distance (version 6.0, release 2, Thomas et al. 2009).

No harbor porpoises were detected in an abundance survey that was conducted concurrently (June–August 2011) in waters between central Virginia and central Florida. This shipboard survey included shelf-break and inner continental slope waters deeper than the 50-m depth contour within the U.S. EEZ. The survey employed the double-platform methodology searching with 25x150 “bigeye” binoculars. A total of 4,445 km of tracklines was surveyed, yielding 290 cetacean sightings.

Table 1. Summary of recent abundance estimates for the Gulf of Maine/Bay of Fundy harbor porpoise (*Phocoena phocoena phocoena*) by month, year, and area covered during each abundance survey and the resulting abundance estimate (Nbest) and coefficient of variation (CV).

Month/Year	Area	Nbest	CV
Jul-Aug 2011	Central Virginia to lower Bay of Fundy	79,883	0.32

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normal distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as

specified by Wade and Angliss (1997). The best estimate of abundance for harbor porpoises is 79,883 (CV=0.32). The minimum population estimate for the Gulf of Maine/Bay of Fundy harbor porpoise is 61,415.

Current Population Trend

A trend analysis has not been conducted for this stock. The statistical power to detect a trend in abundance for this stock is poor due to the relatively imprecise abundance estimates and long survey interval. For example, the power to detect a precipitous decline in abundance (i.e., 50% decrease in 15 years) with estimates of low precision (e.g., CV > 0.30) remains below 80% (alpha = 0.30) unless surveys are conducted on an annual basis (Taylor et al. 2007).

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Several attempts have been made to estimate potential population growth rates. Barlow and Boveng (1991), who used a re-scaled human life table, estimated the upper bound of the annual potential growth rate to be 9.4%. Woodley and Read (1991) used a re-scaled Himalayan tahr life table to estimate a likely annual growth rate of 4%. In an attempt to estimate a potential population growth rate that incorporates many of the uncertainties in survivorship and reproduction, Caswell et al. (1998) used a Monte Carlo method to calculate a probability distribution of growth rates. The median potential annual rate of increase was approximately 10%, with a 90% confidence interval of 3–15%. This analysis underscored the considerable uncertainty that exists regarding the potential rate of increase in this population. Moore and Read (2008) conducted a Bayesian population modeling analysis to estimate the potential population growth of harbor porpoise in the absence of bycatch mortality. Their method used fertility data, in combination with age-at-death data from stranded animals and animals taken in gillnets, and was applied under two scenarios to correct for possible data bias associated with observed bycatch of calves. Demographic parameter estimates were ‘model averaged’ across these scenarios. The Bayesian posterior median estimate for potential natural growth rate was 0.046. This last, most recent, value will be the one used for the purpose of this assessment.

Key uncertainties in the estimate of the maximum net productivity rate for this stock were discussed in Moore and Read (2008), which included the assumption that the age structure is stable, and the lack of data to estimate the probability of survivorship to maximum age. The authors considered the effects of these uncertainties on the estimated potential natural growth rate to be minimal.

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a recovery factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is 61,415. The maximum productivity rate is 0.046. The recovery factor is 0.5 because stock's status relative to OSP is unknown and the CV of the average mortality estimate is less than 0.3 (Wade and Angliss 1997). PBR for the Gulf of Maine/Bay of Fundy harbor porpoise is 706.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

The total annual estimated average human-caused mortality and serious injury is 256 harbor porpoises per year (CV=0.18) from U.S. fisheries using observer data. As the current abundance estimate is only for animals in U.S. waters, Canadian bycatch is not included in the human-caused mortality estimate.

A key uncertainty is the potential that the observer coverage in the Mid-Atlantic gillnet may not be representative of the fishery during all times and places, since the observer coverage was relatively low for some times and areas, 0.02 – 0.10. The effect of this is unknown.

There are no major known sources of unquantifiable human-caused mortality or serious injury for the US waters of the Gulf of Maine/Bay of Fundy population. However, mortality in Canadian Atlantic waters is largely unquantified.

Fishery Information

Detailed U.S. fishery information is reported in Appendix III.

Earlier Interactions

See Appendix V for more information on historical takes.

U.S.

Northeast Sink Gillnet

Harbor porpoise bycatch in the northern Gulf of Maine occurs primarily from June to September, while in the southern Gulf of Maine and south of New England, bycatch occurs from January to May and September to December. Annual bycatch is estimated using ratio estimator techniques that account for the use of pingers (Hatch

and Orphanides 2014, 2015, 2016, Orphanides and Hatch 2017, Orphanides 2019). See Table 2 for bycatch estimates and observed mortality and serious injury for the current 5-year period, and Appendix V for historical bycatch information.

Mid-Atlantic Gillnet

Harbor porpoise bycatch in Mid-Atlantic waters occurs primarily from December to May in waters off New Jersey and less frequently in other waters ranging farther south, from New Jersey to North Carolina. Annual bycatch is estimated using ratio estimator techniques (Hatch and Orphanides 2014, 2015, 2016; Orphanides and Hatch 2017, Orphanides 2019). See Table 2 for bycatch estimates and observed mortality and serious injury for the current 5-year period, and Appendix V for historical bycatch information.

Northeast Bottom Trawl

Since 1989, harbor porpoise mortalities have been observed in the northeast bottom trawl fishery, but many of these were not attributable to this fishery because decomposed animals are presumed to have been dead prior to being taken by the trawl. Those infrequently caught freshly dead harbor porpoises have been caught during January to April on Georges Bank or in the southern Gulf of Maine. Fishery-related bycatch rates were estimated using an annual stratified ratio-estimator (Chavez-Rosales 2018). See Table 2 for bycatch estimates and observed mortality and serious injury for the current 5-year period, and Appendix V for historical bycatch information.

CANADA

No current estimates exist, but harbor porpoise interactions have been documented in the Bay of Fundy sink gillnet fishery and in herring weirs between the years 1998-2001 in the lower Bay of Fundy demersal gillnet fishery (Trippel and Shepherd 2004). That fishery has declined since 2001 and it is assumed bycatch is very small, if any (H. Stone, Department of Fisheries and Oceans Canada, pers. comm.).

Table 2. From observer program data, summary of the incidental mortality of Gulf of Maine/Bay of Fundy harbor porpoise (*Phocoena phocoena phocoena*) by commercial fishery including the years sampled, the type of data used, the annual observer coverage, the mortalities and serious injuries recorded by on-board observers, the estimated annual serious injury and mortality, the estimated CV of the annual mortality, and the mean annual combined mortality (CV in parentheses).

Fishery	Years	Data Type a	Observer Coverage b	Observed Serious Injury	Observed Mortality	Estimated Serious Injury	Estimated Mortality	Estimated Combined Mortality	Estimated CVs	Mean Combined Annual Mortality
Northeast Sink Gillnet	2012	Obs. Data, Trip	0.15	0	34	0	277	277	0.59	221 (.020)
	2013	Logbook, Allocated	0.11	0	20	0	399	399	.33	
	2014	Dealer Data	0.18	0	28	0	128	128	.27	
	2015		0.14	0	23	0	177	177	.28	
	2016		0.10	0	11	0	125	125	.34	
Mid-Atlantic Gillnet	2012	Obs. Data, Weighout	0.02	0	2	0	63	63	0.83	32 (.46)
	2013		0.03	0	1	0	19	19	1.06	
	2014		0.05	0	1	0	22	22	1.03	
	2015		0.06	0	2	0	33	33	1.16	
	2016		0.08	0	2	0	23	23	0.64	
Northeast Bottom Trawl	2012	Obs. Data, Weighout	0.17	0	0	0	0	0	0	3.2 (0.53)
	2013		0.15	0	1	0	7	7	0.98	
	2014		0.17	0	1	0	5.5	5.5	0.86	
	2015		0.19	0	4	0	3.71	3.71	0.49	
	2016		0.12	0	0	0	0	0	0	
TOTAL	-	-	-	-	-	-	-	-	-	256 (0.18)

a Observer data (Obs. Data) are used to measure bycatch rates and the data are collected within the Northeast Fisheries Observer Program. NEFSC collects Weighout (Weighout) landings data that are used as a measure of total effort for the U.S. gillnet fisheries. Mandatory vessel trip report (VTR) (Trip Logbook) data are used to determine the spatial distribution of fishing effort in the northeast sink gillnet fishery.

b Observer coverage for the U.S. Northeast and mid-Atlantic coastal gillnet fisheries is based on tons of fish landed. Northeast bottom trawl

fishery coverages are ratios based on trips.

c Serious injuries were evaluated for the 2012–2016 period and include both at-sea monitor and traditional observer data (Josephson et al. 2019).

Other Mortality

U.S.

There is evidence that harbor porpoises were harvested by natives in Maine and Canada before the 1960s, and the meat was used for human consumption, oil, and fish bait (NMFS 1992). The extent of these past harvests is unknown, though it is believed to have been small. Up until the early 1980s, small kills by native hunters (Passamaquoddy Indians) were reported. It was believed to have nearly stopped (Polacheck 1989) until media reports in September 1997 depicted a Passamaquoddy tribe member dressing out a harbor porpoise. Further articles describing use of porpoise products for food and other purposes were timed to coincide with ongoing legal action in state court.

Recent harbor porpoise strandings on the U.S. Atlantic coast are documented in Table 3 (NOAA National Marine Mammal Health and Stranding Response Database unpublished data, accessed 03 November 2017).

During 2012, 45 harbor porpoises were reported stranded on Atlantic U.S. beaches. Of these, 4 stranding mortalities were reported as having signs of human interaction, one of which was reported to be a fishery interaction.

During 2013, 102 harbor porpoises were reported stranded on Atlantic U.S. beaches. Of these, 9 stranding mortalities were reported as having signs of human interaction, three of which were reported to be fishery interactions.

During 2014, 39 harbor porpoises were reported stranded on Atlantic U.S. beaches. Of these, 5 stranding mortalities were reported as having signs of human interactions, one of which was reported to have been a fishery interaction.

During 2015, 44 harbor porpoises were reported stranded on Atlantic U.S. beaches. Of these, 2 stranding mortalities were reported as having signs of human interactions, neither of which were fishery interactions.

During 2016, 25 harbor porpoises were reported stranded on Atlantic U.S. beaches. Of these, 2 stranding mortalities were reported as having signs of human interactions, one of which was reported to have been a fishery interaction.

Stranding data probably underestimate the extent of fishery-related mortality and serious injury because all of the marine mammals that die or are seriously injured may not wash ashore, nor will all of those that do wash ashore necessarily show signs of entanglement or other fishery-interaction. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of fishery interaction.

Table 3. Harbor porpoise (*Phocoena phocoena phocoena*) reported strandings along the U.S. and Canadian Atlantic coast, 2012-2016.

Area	2012	2013	2014	2015	2016	Total
Main ^{ea, d, e}	7	7	5	2	5	26
New Hampshire	3	1	1	0	1	6
Massachusetts ^{a, c, d, e}	25	40	22	18	8	113
Rhode Island ^{f, g}	0	3	0	2	2	7
Connecticut ^d	0	1	0	0	0	1
New York ^{b, d}	3	15	1	3	1	23
New Jersey ^{a, d, e}	2	8	4	2	5	21
Delaware	0	2	0	0	0	2
Maryland	1	3	0	0	0	4

Area	2012	2013	2014	2015	2016	Total
Virginia ^{c, g}	2	15	3	3	2	25
North Carolina ^f	2	7	11	14	1	62
TOTAL U.S.	45	102	39	44	25	263
Nova Scotia/Prince Edward Island ^h	6	21	9	13	16	62
Newfoundland and New Brunswick	0	3	0	2	0	5
GRAND TOTAL	51	126	48	59	16	461

a. One Maine animal was taken to rehab in 2012. Three Massachusetts live strandings were taken to rehab in 2013 and 1 Maine animal was released alive. In 2016, one animal in Maine and one animal in New Jersey were responded to and released alive.

b. One of the 2012 New York strandings classified as human interaction due to interaction with marine debris.

c. Four HI cases in 2012: one of these was a fishery interaction (Massachusetts).

d. Ten total HI cases in 2013 (MA-3, ME-2, NY-3, NJ-1, CT-1), including one released alive (ME). Three of these were considered fishery interactions, including one entangled in gear in Maine.

e. Five total HI cases in 2014: 2 in Maine, 1 each in Massachusetts, New Jersey and Virginia. The Virginia case was recorded as a fishery interaction.

f. Two HI cases in 2015: 1 in Rhode Island and 1 in North Carolina

g. Two HI cases in 2016: 1 in Rhode Island and 1 in Virginia. The Virginia case was coded as a fishery interaction.

h. Data supplied by Nova Scotia Marine Animal Response Society (pers. comm.). One of the 2012 animals trapped in mackerel net. Not included in count for 2014 are at least 8 animals released alive from weirs. One of the 2015 animals a suspected fishery interaction.

i. (Ledwell and Huntington 2012a, 2012b, 2013, 2014, 2015, 2017).

CANADA

Whales and dolphins stranded on the coast of Nova Scotia, New Brunswick and Prince Edward Island are recorded by the Marine Animal Response Society and the Nova Scotia Stranding Network. See Table 3 for details.

Harbor porpoises stranded on the coasts of Newfoundland and Labrador are reported by the Newfoundland and Labrador Whale Release and Strandings Program (Ledwell and Huntington 2012a, 2012b, 2013; 2014; Table 3).

HABITAT ISSUES

Harbor porpoise are mostly found in nearshore areas and inland waters, including bays, tidal areas, and river mouths. As a result, in addition to fishery bycatch, harbor porpoise are vulnerable to contaminants, such as PCBs (Hall et al. 2006), ship traffic (Oakley et al. 2017; Terhune 2015) and physical modifications resulting from urban and industrial development activities such as construction of docks and other over-water structures, dredging (Todd et al. 2015), installation of offshore windfarms (Carstensen et al. 2006; Dähne et al. 2013; Benjamins et al. 2017), seismic surveys and noise (Lucke et al. 2009).

STATUS OF STOCK

Harbor porpoise in the Gulf of Maine/Bay of Fundy are not listed as threatened or endangered under the Endangered Species Act, and this stock is not considered strategic under the MMPA. The total U.S. fishery-related mortality and serious injury for this stock is not less than 10% of the calculated PBR and, therefore, cannot be considered to be insignificant and approaching zero mortality and serious injury rate. The status of harbor porpoises, relative to OSP, in the U.S. Atlantic EEZ is unknown. Population trends for this species have not been investigated.

Based on the low levels of uncertainties described in the above sections, it expected these uncertainties will have little effect on the designation of the status of this stock.

REFERENCES CITED

- Barlow, J. and P. Boveng 1991. Modeling age-specific mortality for marine mammal populations. *Mar. Mamm. Sci.* 7: 50–65.
- Benjamins, S., N. van Geel, G. Hastie, J. Elliott and B. Wilson. 2017. Harbour porpoise distribution can vary at small spatiotemporal scales in energetic habitats. *Deep-Sea Res. II.* <http://dx.doi.org/10.1016/j.dsr2.2016.07.002i>

- Carstensen J, O.D. Henriksen and J. Teilmann. Impacts of offshore wind farm construction on harbour porpoises: acoustic monitoring of echolocation activity using porpoise detectors (T-PODs). *Mar Ecol Prog Ser.* 2006;321:295–308.
- Caswell, H., S. Brault, A.J. Read and T.D. Smith 1998. Harbor porpoise and fisheries: An uncertainty analysis of incidental mortality. *Ecol. Appl.* 8(4): 1226–1238.
- Chavez-Rosales, S., M.C. Lyssikatos, and J. Hatch. 2018. Estimates of cetacean and pinniped bycatch in northeast and mid-Atlantic bottom trawl fisheries, 2012-2016. NOAA Tech. Memo NMFS-NE-250. 29pp.
- Dähne, M. A. Gilles, K. Lucke, V. Peschko, S. Adler, K. Krugel, J. Sundermeyer and U. Siebert. 2013. Effects of pile-driving on harbour porpoises (*Phocoena phocoena*) at the first offshore wind farm in Germany. *Environ. Res. Lett.* 8: 1–15.
- Gaskin, D.E. 1977. Harbour porpoise, *Phocoena phocoena* (L.), in the western approaches to the Bay of Fundy 1969-75. *Rep. Int. Whal. Comm.* 27: 487–492.
- Gaskin, D.E. 1984. The harbor porpoise *Phocoena phocoena* (L.): Regional populations, status, and information on direct and indirect catches. *Rep. Int. Whal. Comm.* 34: 569–586.
- Gaskin, D.E. 1992. The status of the harbour porpoise. *Can. Field-Nat.* 106: 36–54.
- Hall, A.J., K. Hugunin, R. Deaville, R.J. Law, C.R. Allchin and P.D. Jepson 2006. The risk of infection from polychlorinated biphenyl exposure in the harbor porpoise (*Phocoena phocoena*): A case-control approach. *Environmental Health Perspectives* 114(5): 704–711.
- Hatch, J.M. and C.D. Orphanides 2016. Estimates of cetacean and pinniped bycatch in the 2014 New England sink and mid-Atlantic gillnet fisheries. *Northeast Fish Sci Cent Ref Doc* 16-05. 22 pp.
- Hatch, J.M. and C.D. Orphanides 2015. Estimates of cetacean and pinniped bycatch in the 2013 New England sink and mid-Atlantic gillnet fisheries. *Northeast Fish Sci Cent Ref Doc.* 15–15; 26 p. Available at <http://www.nefsc.noaa.gov/publications/> doi:10.7289/V5HD7SNK.
- Hatch, J.M. and C.D. Orphanides 2014. Estimates of cetacean and pinniped bycatch in the 2012 New England sink and mid-Atlantic gillnet fisheries. *Northeast Fish Sci Cent Ref Doc.* 14-02. 20 pp.
- Hiltunen, K.H. 2006. Mixed-stock analysis of harbor porpoises (*Phocoena phocoena*) along the U.S. mid-Atlantic coast using microsatellite DNA markers. MS thesis. The College of Charleston, Charleston, SC. 92 pp.
- Johnston, D.W. 1995. Spatial and temporal differences in heavy metal concentrations in the tissues of harbour porpoises (*Phocoena phocoena* L.) from the western North Atlantic. M.S. thesis. University of Guelph, Guelph, Ontario, Canada. 152 pp.
- Josephson, E., F. Wenzel and M.C. Lyssikatos. 2019. [Serious injury determinations for small cetaceans and pinnipeds caught in commercial fisheries off the northeast U.S. coast, 2012-2016](#). *Northeast Fish. Sci. Cent. Ref. Doc.* 19-05, 27pp.
- Kraus, S.D., J.H. Prescott and G.S. Stone 1983. Harbor porpoise, *Phocoena phocoena*, in the U.S. coastal waters off the Gulf of Maine: a survey to determine seasonal distribution and abundance. *NMFS. NA82FAC00027* 22 pp.
- Kraus, S.D., A.J. Read, A. Solow, K. Baldwin, T. Spradlin, E. Anderson and J. Williamson 1997. Acoustic alarms reduce porpoise mortality. *Nature* 388(6642): 525.
- Laake, J.L., and D.L. Borchers 2004. Methods for incomplete detection at distance zero, In: *Advanced distance sampling*, edited by S. T. Buckland, D. R. Andersen, K. P. Burnham, J. L. Laake, and L. Thomas, pp. 108–189, Oxford University Press, New York.
- Ledwell, W. and J. Huntington 2012a. Incidental entrapments of cetaceans and leatherback sea turtles in fishing gear reported during 2011-2012 and a summary of the Whale Release and Stranding Group activities. A preliminary report to Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 17 pp.
- Ledwell, W. and J. Huntington 2012b. Incidental entrapments in fishing gear and stranding reported to and responded to by the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the Whale Release and Strandings program during 2012. Report to Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 18 pp.
- Ledwell, W., J. Huntington and E. Sacrey 2013. Incidental entrapments in fishing gear and stranding reported to and responded to by the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the Whale Release and Strandings program during 2013. Report to the Department of Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 19 pp.
- Ledwell, W. and J. Huntington 2014. Incidental entrapments and entanglements of cetaceans and leatherback sea turtles, strandings, ice entrapments reported to the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the Whale Release and Strandings program during 2014. Report to the Department of Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 23 pp.

- Ledwell, W. and J. Huntington 2015. Incidental entrapments and entanglements of cetaceans and leatherback sea turtles, strandings, ice entrapments reported to the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the Whale Release and Strandings program during 2015. Report to the Department of Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 22 pp.
- Ledwell, W. and J. Huntington 2017. Incidental entrapments and entanglements of cetaceans and leatherback sea turtles and strandings and harassments reported to the Whale Release and Strandings Group in Newfoundland and Labrador and a summary of the Whale Release and Strandings program during 2016. Report to the Department of Fisheries and Oceans Canada, St. John's, Newfoundland, Canada. 23 pp.
- Lucke K., U. Siebert, P.A. Lepper and M-A. Blanchet. 2009. Temporary shift in masking hearing thresholds in a harbor porpoise (*Phocoena phocoena*) after exposure to seismic airgun stimuli. *J. Acoust. Soc. Am.* 125(6):4060–4070.
- Lyssikatos, M. 2015. Estimates of Cetacean and Pinniped Bycatch in Northeast and mid-Atlantic bottom trawl fisheries, 2008-2013. Northeast Fish Sci Cent Ref Doc. 15-19. 20 p. Available at <http://www.nefsc.noaa.gov/publications/>.
- Moore, J.E. and A.J. Read 2008. A Bayesian uncertainty analysis of cetacean demography and bycatch mortality using age-at-death data. *Ecol. Appl.* 18(8): 1914–1931.
- NMFS 1992. Harbor porpoise in Eastern North America: Status and Research Needs. Results of a scientific workshop held May 5-8, 1992 at NEFSC, Woods Hole, MA, USA. Northeast Fish. Sci. Cent. Ref. Doc., 92-06. National Marine Fisheries Service. 28 pp.
- Oakley, J.A., A.T. Williams and T. Thomas. 2017. Reactions of harbor porpoise (*Phocoena phocoena*) to vessel traffic in the coastal waters of South West Wales, UK. *Ocean & Coastal Management* 138:158–169.
- Orphanides, C.D. 2019. Estimates of cetacean and pinniped bycatch during 2016 in the New England and Mid-Atlantic Sink Gillnet fisheries. Northeast Fish Sci Cent Ref Doc. 19-04. 12 pp.
- Orphanides, C.D. and Hatch, J. 2017. Estimates of cetacean and pinniped bycatch during 2015 in the New England and Mid-Atlantic Sink Gillnet fisheries. Northeast Fish Sci Cent Ref Doc. 17-18. 21pp.
- Orphanides, C.D. 2013. Estimates of cetacean and pinniped bycatch during 2010 and 2011 in the New England Sink Gillnet fishery, Mid-Atlantic Gillnet fishery, and two NMFS gillnet experiments. Northeast Fish Sci Cent Ref Doc. 13-13 38 pp. Available at: <http://nefsc.noaa.gov/publications/crd/crd1313/>.
- Palka, D. 1995. Influences on spatial patterns of Gulf of Maine harbor porpoises. Pages 69–75 in: A. S. Blix, L. Walloe and O. Ulltang, (eds.) *Whales, Seals, Fish and Man*. Elsevier Science. Amsterdam.
- Palka, D.L., A.J. Read, A.J. Westgate and D.W. Johnston 1996. Summary of current knowledge of harbour porpoises in US and Canadian Atlantic waters. *Rep. Int. Whal. Comm.* 46: 559–565.
- Palka, D. 2000. Abundance of the Gulf of Maine/Bay of Fundy harbor porpoise based on shipboard and aerial surveys during 1999. Northeast Fish. Sci. Cent. Ref. Doc. 00-07. 29 pp. <http://www.nefsc.noaa.gov/psb/pubs/palkalabref00-07.pdf>
- Palka, D.L. 2012. Cetacean abundance estimates in US northwestern Atlantic Ocean waters from summer 2011 line transect survey. Northeast Fish. Sci. Cent. Ref. Doc. 12-29. 37 pp. <http://www.nefsc.noaa.gov/nefsc/publications/crd/crd1229/>
- Polacheck, T. 1989. Harbor porpoises and the gillnet fishery. *Oceanus* 32(1): 63-70.
- Read, A.J. and A.A. Hohn 1995. Life in the fast lane: the life history of harbour porpoises from the Gulf of Maine. *Mar. Mamm. Sci.* 11(4): 423–440.
- Read, A.J. and A.J. Westgate 1997. Monitoring the movements of harbour porpoises (*Phocoena phocoena*) with satellite telemetry. *Ma. Biol.* 130: 315–22.
- Rosel, P.E., S.C. France, J.Y. Wang and T.D. Kocher 1999a. Genetic structure of harbour porpoise *Phocoena phocoena* populations in the northwest Atlantic based on mitochondrial and nuclear markers. *Mol. Ecol.* 8: S41-S54.
- Rosel, P.E., R. Tiedemann and M. Walton 1999b. Genetic evidence for limited trans-Atlantic movements of the harbor porpoise *Phocoena phocoena*. *Mar. Biol.* 133: 583–591.
- Taylor, B.L., M. Martinez, T. Gerrodette, J. Barlow and Y.N. Hrovat. 2007. Lessons from monitoring trends in abundance in marine mammals. *Mar. Mamm. Sci.* 23(1): 157–175.
- Terhune, J.M. 2015. Harbour porpoise presence near oil tankers. *J. Canadian Ac.* 43(3) 2pp.
- Thomas L., J.L. Laake, E. Rexstad, S. Strindberg, F.F.C. Marques, S.T. Buckland, D.L. Borchers, D.R. Anderson, K.P. Burnham, M.L. Burt, S.L. Hedley, J.H. Pollard, J.R.B. Bishop and T.A. Marques. 2009. Distance 6.0. Release 2. [Internet]. University of St. Andrews (UK): Research Unit for Wildlife Population Assessment. Available from: <http://distancesampling.org/Distance/>

- Todd, V.L.G., I.B. Todd, J.C. Gardiner, E.C.N. Morrin, N.A. MacPherson, N.A. DiMarzio and F. Thomsen. 2015. A review of impacts of marine dredging activities on marine mammals. *ICES J. of Mar. Sci.* 72(2): 328-340.
- Trippel, E.A. and T.D. Shepherd 2004. By-catch of harbour porpoise (*Phocoena phocoena*) in the Lower Bay of Fundy gillnet fishery from 1998-2001. Department of Fisheries and Oceans. Ottawa, Ontario. DFO Research Document 2004/2521 iv + 33 pp. http://www.fmap.ca/ramweb/papers-total/Trippel_Shepherd_2004.pdf
- Wade, P.R. and R.P. Angliss 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. NOAA Tech. Memo. NMFS-OPR-12. 93 pp.
- Wang, J.Y., D.E. Gaskin and B.N. White 1996. Mitochondrial DNA analysis of harbour porpoise, *Phocoena phocoena*, subpopulations in North American waters. *Can. J. Fish. Aquat. Sci* 53: 1632-45.
- Westgate, A.J., D.C.G. Muir, D.E. Gaskin and M.C.S. Kingsley 1997. Concentrations and accumulation patterns of organochlorine contaminants in the blubber of harbour porpoises, *Phocoena phocoena*, from the coast of Newfoundland, the Gulf of St. Lawrence and the Bay of Fundy/Gulf of Maine. *Envir. Pollut.* 95: 105-119.
- Westgate, A.J., A.J. Read, T.M. Cox, T.D. Schofield, B.R. Whitaker and K.E. Anderson 1998. Monitoring a rehabilitated harbor porpoise using satellite telemetry. *Mar. Mamm. Sci.* 14(3): 599-604.
- Westgate, A.J. and K.A. Tolley 1999. Geographical differences in organochlorine contaminants in harbour porpoises *Phocoena phocoena* from the western North Atlantic. *Mar. Ecol. Prog. Ser.* 177: 255-268.
- Wingfield J.E., M. O'Brien, V. Lyubchich, J.J. Roberts, P.N. Halpin, A.N. Rice and H. Bailey. (2017) Year-round spatiotemporal distribution of harbor porpoises within and around the Maryland wind energy area. *PLoS ONE* 12(5): e0176653. <https://doi.org/10.1371/journal.pone.0176653>
- Woodley, T.H. and A.J. Read 1991. Potential rates of increase of a harbor porpoise (*Phocoena phocoena*) population subjected to incidental mortality in commercial fisheries. *Can. J. Fish. Aquat. Sci* 48: 2429-35.