


## **HUMPBACK WHALE (*Megaptera novaeangliae*): Western North Pacific Stock**

### **STOCK DEFINITION AND GEOGRAPHIC RANGE**

The humpback whale is distributed worldwide in all ocean basins, though it is less common in Arctic waters. In winter, most humpback whales occur in the temperate and tropical waters of the North and South Hemispheres (from 10°-23° latitude). Humpback whales in the North Pacific are seasonal migrants that feed on zooplankton and small schooling fishes in the cool, coastal waters of the western United States, western Canada, and the Russian Far East (NMFS 1991). The historic feeding range of humpback whales in the North Pacific encompassed coastal and inland waters around the Pacific rim from Point Conception, California, north to the Gulf of Alaska and the Bering Sea, and west along the Aleutian Islands to the Kamchatka Peninsula and into the Sea of Okhotsk (Nemoto 1957, Tomlin 1967, Johnson and Wolman 1984). Humpback whales have been known to enter the Chukchi Sea (Johnson and Wolman 1984). The humpback whale population in much of this range was considerably reduced as a result of intensive commercial exploitation during this century.



**Figure 31.** Approximate distribution of humpback whales in the western North Pacific (shaded area). Feeding and wintering areas are presented above (see text). See Figure 32 for humpback whale distribution in the eastern North Pacific.

Aerial, vessel, and photo-identification surveys and genetic analyses indicate that within the U. S. Exclusive Economic Zone (EEZ) there are at least three relatively separate populations that migrate between their respective summer/fall feeding areas to winter/spring calving and mating areas (Calambokidis et al. 1997, Baker et al. 1998, Figs. 31 and 32): 1) winter/spring populations in coastal Central America and Mexico which migrate to the coast of California to southern British Columbia in summer/fall (Calambokidis et al. 1989, Steiger et al. 1991, Calambokidis et al. 1993) - referred to as the California/Oregon/Washington and Mexico stock; 2) winter/spring populations of the Hawaiian Islands which migrate to northern British Columbia/Southeast Alaska and Prince William Sound west to Kodiak (Baker et al. 1990, Perry et al. 1990, Calambokidis et al. 1997) - referred to as the Central North Pacific stock; and 3) winter/spring populations of Japan which, based on Discovery Tag information, probably migrate to waters west of the Kodiak Archipelago (the Bering Sea and Aleutian Islands) in summer/fall (Berzin and Rovnin 1966, Nishiwaki 1966, Darling 1991) - referred to as the Western North Pacific stock. Winter/spring populations of humpback whales also occur in Mexico's offshore islands. The migratory destination of these whales is not well known (Calambokidis et al. 1993, Calambokidis et al. 1997). Some recent exchange between winter/spring areas has been documented (Darling and McSweeney 1985, Baker et al. 1986,

Darling and Cerchio 1993), as well as movement between Japan and British Columbia, and Japan and the Kodiak Archipelago (Darling et al. 1996, Calambokidis et al. 1997).

Currently, there are insufficient data to apply the Dizon et al.(1992) phylogeographic approach to classify population structure in humpback whales. Until further information becomes available, 3 management units of humpback whales (as described above) are recognized within the U.S. EEZ of the North Pacific: one in the Eastern North Pacific (the California/Oregon/Washington - Mexico stock), one in the Central North Pacific, and one in the Western North Pacific. The California/Oregon/Washington - Mexico humpback whale stock is reported separately in the Stock Assessment Reports for the Pacific Region.

The feeding areas for the Western North Pacific humpback whale stock are largely unknown. There has been little to no effort to photo-identify individual humpback whales in the North Pacific waters west of the Kodiak Archipelago. As a result, none of the whales identified off Japan have been resighted in the historical feeding areas of the stock (Bering Sea and Aleutian Islands). Individuals identified off Japan, however, have been resighted in the eastern North Pacific (Calambokidis et al. 1997). This may indicate that the Western North Pacific humpback whale stock did not exclusively use the feeding areas in the western Pacific, or, perhaps, a shift in the migratory destination of this stock has occurred. Thus, some unknown fraction of whales from the wintering grounds off Japan spend their summers feeding in areas typically utilized by whales from the Central North Pacific stock.

### **POPULATION SIZE**

The abundance estimate of humpback whales in the North Pacific is based on data collected by nine independent research groups that conducted photo-identification studies of humpback whales in the three wintering areas (Mexico, Hawaii, and Japan). Photographs taken between 1991 and 1993 were used to estimate abundance because samples throughout the entire North Pacific were the largest and most complete during this period. Using Darroch's (1961) method, which utilizes only data from wintering areas (in this case data provided by two Japanese research groups), and averaging the 1991-92, 1992-93, and 1991-93 winter release-recovery information results in an abundance estimate of 394 (CV=0.084) for the Western North Pacific humpback whale stock (Calambokidis et al. 1997).

A vessel survey conducted in August of 1994 covered 2,050 nautical miles of trackline south of the Aleutian Islands encountered humpback whales in scattered aggregations (57 sightings) throughout the study area (Forney and Brownell 1996). It is unknown whether the humpback whales encountered during this survey belonged to the Western or Central North Pacific stock.

There are no reliable estimates for the abundance of humpback whales at feeding areas for this stock because the specific feeding areas are largely unknown.

### **Minimum Population Estimate**

The minimum population estimate ( $N_{MIN}$ ) for this stock is calculated according to Equation 1 from the PBR Guidelines (Wade and Angliss 1997):  $N_{MIN} = N/\exp(0.842 \times [\ln(1+[CV(N)]^2)]^{1/2})$ . Using the population estimate (N) of 394 and its associated CV(N) of 0.084,  $N_{MIN}$  for this humpback whale stock is 367.

### **Current Population Trend**

Reliable information on trends in abundance for the Western North Pacific humpback whale stock are currently not available.

### **CURRENT AND MAXIMUM NET PRODUCTIVITY RATES**

Utilizing a birth-interval model, Barlow and Clapham (1997) have estimated a population growth rate of 6.5% (SE=1.2%) for the well-studied humpback whale population in the Gulf of Maine. However, there are no estimates of the growth rate of humpback whale populations in the North Pacific (Best 1993). Hence, until additional data become available from this or other North Pacific humpback whale stocks, it is recommended that the cetacean maximum net productivity rate ( $R_{MAX}$ ) of 4% be employed for this stock (Wade and Angliss 1997).

### **POTENTIAL BIOLOGICAL REMOVAL**

Under the 1994 re-authorized Marine Mammal Protection Act (MMPA), the potential biological removal (PBR) is defined as the product of the minimum population estimate, one-half the maximum theoretical net productivity rate, and a recovery factor:  $PBR = N_{MIN} \times 0.5R_{MAX} \times F_R$ . The recovery factor ( $F_R$ ) for this stock is 0.1, the value for cetacean stocks

listed as endangered under the Endangered Species Act (Wade and Angliss 1997). Thus, for the Western North Pacific stock of humpback whale, PBR = 0.7 animals ( $367 \times 0.02 \times 0.1$ ).

## ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

### Fisheries Information

Six different commercial fisheries operating in Alaska waters within the range of this stock were monitored for incidental take by fishery observers during 1990-98: Bering Sea/Aleutian Islands groundfish trawl, longline, and pot fisheries, and Gulf of Alaska groundfish trawl, longline, and pot fisheries. One humpback whale mortality was observed in the Bering Sea/Aleutian Islands groundfish trawl fishery in 1998. Average annual mortality from observed fisheries was 0.2 humpbacks from this stock (Table 25). Note, however, that the stock identification is uncertain and the mortality may have been attributable to the central stock of humpback whales. Thus, this mortality is assigned to both the central and western stocks.

An additional source of information on the number of humpback whales killed or injured incidental to commercial fishery operations is the self-reported fisheries information required of vessel operators by the MMPA. During the period between 1990 and 1998, there were no fisher self-reports of humpback whale injuries or mortalities from interactions with commercial fishing gear in any Alaska fishery within the presumed range of the Western North Pacific humpback whale stock. Logbook data are available for part of 1989-1994, after which incidental mortality reporting requirements were modified. Under the new system, logbooks are no longer required; instead, fishers provide self-reports. Data for the 1994-95 phase-in period is fragmentary. After 1995, the level of reporting dropped dramatically, such that the records are considered incomplete and estimates of mortality based on them represent minimums (see Appendix 4 for details).

Strandings of humpback whales entangled in fishing gear or with injuries caused by interactions with gear are another source of mortality data. The only fishery-related humpback stranding in an area thought to be occupied by animals from this stock was reported by a U. S. Coast Guard vessel in late June 1997 operating near the Bering Strait. The whale was found floating dead entangled in netting and trailing orange buoys (National Marine Mammal Laboratory, Platforms of Opportunity Program, unpubl. data, 7600 Sand Point Way NE, Seattle, WA 98115). With the given data it is not possible to determine which fishery (or even which country) caused the mortality. Note, that this mortality has been attributed the Western North Pacific stock, but without a tissue sample (for genetic analysis) or a photograph (for matching to known Japanese animals) it is not possible to for certain (i.e., it may have belonged to the Central North Pacific stock). Averaging this mortality over the 5-year period 1994-98 results in an estimated annual mortality of 0.2 humpback whales from this stock. This estimate is considered a minimum because not all entangled animals strand and not all stranded animals are found, or reported.

**Table 25.** Summary of incidental mortality of humpback whales (Western North Pacific stock) due to commercial fisheries from 1990 through 1998 and calculation of the mean annual mortality rate. Mean annual mortality in brackets represents a minimum estimate. For a particular fishery, the most recent 5 years of available data are used in the mortality calculation when more than 5 years of data are provided. n/a indicates that data are not available.

Fishery name	Years	Data type	Range of observer coverage	Observed mortality (in given yrs.)	Estimated mortality (in given yrs.)	Mean annual mortality
Bering Sea/Aleutian Is. (BSA) groundfish trawl	90-98	obs data	53-74%	0, 0, 0, 0, 0, 0, 0, 0, 0	0, 0, 0, 0, 0, 0, 0, 0, 1	0.2 (CV=1.0)
Observer program total						0
				<b>Reported mortalities</b>		
unknown fishery (Bering Sea)	94-98	strand data	n/a	0, 0, 0, 1, 0	\$0.2	[\$0.2]

Fishery name	Years	Data type	Range of observer coverage	Observed mortality (in given yrs.)	Estimated mortality (in given yrs.)	Mean annual mortality
Minimum total annual mortality						[\$0.4]

The estimated annual mortality rate incidental to commercial fisheries is 0.4 (0.2 from observed fisheries plus 0.2 from the stranding data) whales per year from this stock. However, this estimate is considered a minimum because there are no data concerning fishery-related mortalities in Japanese, Russian, or international waters. In addition, there is a small probability that fishery interactions discussed in the assessment for the Central North Pacific stock may have involved animals from this stock because the only known matches to feeding areas come from areas typically used by the Central North Pacific stock.

#### Subsistence/Native Harvest Information

Subsistence hunters in Alaska and Russia have not been reported to take humpback whales from this stock.

#### HISTORIC WHALING

The number of humpback whales in the North Pacific may have numbered approximately 15,000 individuals prior to exploitation (Rice 1978). Intensive commercial whaling removed more than 28,000 animals from the North Pacific during the 20th century (Rice 1978). This mortality estimate likely underestimates the actual kill as a result of under-reporting of the Soviet catches (Yablokov 1994).

#### STATUS OF STOCK

As the estimated annual mortality rate (0.4) is considered a minimum, it is unclear whether the level of human-caused mortality and serious injury exceeds the PBR (0.7). At least one of the mortalities occurred in a U. S. fishery; therefore, the estimated fishery mortality and serious injury rate exceeds 10% of the PBR (0.07). The rate cannot be considered insignificant and approaching zero. The humpback whale is listed as “endangered” under the Endangered Species Act, and therefore designated as “depleted” under the MMPA. As a result, the Western North Pacific humpback whale stock is classified as a strategic stock. Reliable population trend data and the status of this stock relative to its Optimum Sustainable Population size are currently unknown. Noise pollution from the U. S. Navy’s Low Frequency Active sonar program and other anthropogenic sources (i.e., shipping) is a potential concern as to the health of this stock.

#### CITATIONS

- Baker, C. S., L. M. Herman, A. Perry, W. S. Lawton, J. M. Straley, A. A. Wolman, G. D. Kaufman, H. E. Winn, J. D. Hall, J. M. Reinke, and J. Ostman. 1986. Migratory movement and population structure of humpback whales (*Megaptera novaeangliae*) in the central and eastern North Pacific. *Mar. Ecol. Prog. Ser.* 31:105-119.
- Baker, C. S., S. R. Palumbi, R. H. Lambertsen, M. T. Weinrich, J. Calambokidis, and S. J. O’Brien. 1990. Influence of seasonal migration on geographic distribution of mitochondrial DNA haplotypes in humpback whales. *Nature* 344:238-240.
- Baker, C. S., L. Medrano-Gonzalez, J. Calambokidis, A. Perry, F. Pichler, H. Rosenbaum, J. M. Straley, J. Urban-Ramirez, M. Yamaguchi, and O. von Ziegeler. 1998. Population structure of nuclear and mitochondrial DNA variation among humpback whales in the North Pacific. *Mol. Ecol.* 7:695-707.
- Barlow, J., and P. J. Clapham. 1997. A new birth-interval approach to estimating demographic parameters of humpback whales. *Ecol.* 78(2):535-546.
- Berzin, A. A., and A. A. Rovnin. 1966. The distribution and migrations of whales in the northeastern part of the Pacific, Chukchi and Bering Seas. *Izvestiya Tikhookeanskogo Nauchno-Issledovatel’skogo Institut Rybnogo Khozyaistva I Okeanografii* 58:179-207. (Translated by Bureau of Commercial Fisheries, U. S. Fish and Wildlife Service, Seattle, 1968, pp. 103-136. *In* K. I. Panin (ed.), *Soviet Research on Marine Mammals of the Far East.*)
- Best, P. B. 1993. Increase rates in severely depleted stocks of baleen whales. *ICES J. Mar. Sci.* 50:169-186.

- Calambokidis, J., G. H. Steiger, J. C. Cubbage, K. C. Balcomb III, and P. Bloedel. 1989. Biology of humpback whales in the Gulf of the Farallones. Report to Gulf of the Farallones National Marine Sanctuary, San Francisco, CA by Cascadia Research Collective, 218½ West Fourth Avenue, Olympia, WA. 93 pp.
- Calambokidis, J., G. H. Steiger, and J. R. Evenson. 1993. Photographic identification and abundance estimates of humpback and blue whales off California in 1991-92. Final Contract Report 50ABNF100137 to Southwest Fisheries Science Center, P.O. Box 271, La Jolla, CA 92038. 67 pp.
- Calambokidis, J., G. H. Steiger, J. M. Straley, T. Quinn, L. M. Herman, S. Cerchio, D. R. Salden, M. Yamaguchi, F. Sato, J. R. Urban, J. Jacobson, O. Von Zeigesar, K. C. Balcomb, C. M. Gabriele, M. E. Dahlheim, N. Higashi, S. Uchida, J. K. B. Ford, Y. Miyamura, P. Ladrón de Guevara, S. A. Mizroch, L. Schlender, and K. Rasmussen. 1997. Abundance and population structure of humpback whales in the North Pacific basin. Final Contract Report 50ABNF500113 to Southwest Fisheries Science Center, P.O. Box 271, La Jolla, CA 92038. 72 pp.
- Darling, J. D. 1991. Humpback whales in Japanese waters. Ogasawara and Okinawa. Fluke identification catalog 1987-1990. Final Contract Report, World Wide Fund for Nature, Japan. 22 pp.
- Darling, J. D., and S. Cerchio. 1993. Movement of a humpback whale (*Megaptera novaeangliae*) between Japan and Hawaii. *Mar. Mammal Sci.* 1:84-89.
- Darling, J. D., and D. J. McSweeney. 1985. Observations on the migrations of North Pacific humpback whales (*Megaptera novaeangliae*). *Can. J. Zool.* 63:308-314.
- Darling, J. D., J. Calambokidis, J., K. C. Balcomb, P. Bloedel, K. Flynn, A. Mochizuki, K. Mori, F. Sato, and M. Yamaguchi. 1996. Movement of a humpback whale (*Megaptera novaeangliae*) from Japan to British Columbia and return. *Mar. Mammal Sci.* 12(2):281-287.
- Darroch, J. N. 1961. The two-sample capture-recapture census when tagging and sampling are stratified. *Biometrika* 48:241-260.
- Dizon, A. E., C. Lockyer, W. F. Perrin, D. P. DeMaster, and J. Sisson. 1992. Rethinking the stock concept: a phylogeographic approach. *Conserv. Biol.* 6:24-36.
- Forney, K. A., and R. L. Brownell. 1996. Preliminary report of the 1994 Aleutian Island marine mammal survey. Unpubl. doc. submitted to Int. Whal. Comm. (SC/48/O 11). 15 pp.
- Johnson, J. H., and A. A. Wolman. 1984. The humpback whale, *Megaptera novaeangliae*. *Mar. Fish. Rev.* 46(4):30-37.
- National Marine Fisheries Service. 1991. Recovery plan for the humpback whale (*Megaptera novaeangliae*). Prepared by the humpback recovery team for the National Marine Fisheries Service, Silver Spring, Maryland. 105 pp.
- Nemoto, T. 1957. Foods of baleen whales in the northern Pacific. *Sci. Rep. Whales Res. Inst. Tokyo* 12:33-89.
- Nishiwaki, M. 1966. Distribution and migration of the larger cetaceans in the North Pacific as shown by Japanese whaling results. Pp. 172-191, *In* K. S. Norris (ed.), *Whales, Dolphins and Porpoises*, University of California Press, Berkeley, CA.
- Perry, A., C. S. Baker, and L. M. Herman. 1990. Population characteristics of individually identified humpback whales in the central and eastern North Pacific: a summary and critique. *Rep. Int. Whal. Comm. (Special Issue 12)*:307-317.
- Rice, D. W. 1978. The humpback whale in the North Pacific: distribution, exploitation and numbers. Appendix 4. Pp. 29-44, *In* K. S. Norris and R. R. Reeves (eds.), *Report on a workshop on problems related to humpback whales (Megaptera novaeangliae) in Hawaii*. U.S. Dep. Commer., Nat. Tech. Info. Serv. PB-280 794. Springfield, VA.
- Steiger, G. H., J. Calambokidis, R. Sears, K. C. Balcomb, and J. C. Cubbage. 1991. Movement of humpback whales between California and Costa Rica. *Mar. Mammal Sci.* 7:306-310.
- Tomlin, A. G. 1967. Mammals of the USSR and adjacent countries. vol. 9, Cetacea. *Israel Program Sci. Transl. No. 1124*, Natl. Tech. Info. Serv. TT 65-50086. Springfield, VA. 717 pp. (Translation of Russian text published in 1957).
- Wade, P. R., and R. Angliss. 1997. Guidelines for assessing marine mammal stocks: report of the GAMMS workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Yablokov, A. V. 1994. Validity of whaling data. *Nature* 367:108.